

ANNUAL INDEX

VOLUME 3

1968

AUTHORS

A

- Alam, A. U., J. R. Couch and C. R. Creger. Short communication re fatty acid composition of the xanthophyll esters of *Tagetes erecta* petals, 183
 Anderson, Carl E. See Edward O. Oswald
 Ansari, M. N. A. See N. Nicolaides
 Argoudelis, C. J. and E. G. Perkins. Short communication re determination of double bond position in mono-unsaturated fatty acids using combination gas chromatography mass spectrometry, 379

B

- Baczynski, E. See D. L. Turner
 Baer, Erich. Short communication re differentiation of nitrogenous phospholipids by infrared spectroscopy, 334
 Baker, Grace L. See Charles L. Sloan
 Baldwin, Jerry and W. E. Cornatzer. Liver lipids during development, 361
 Barclay, A. S. See Roger W. Miller
 Belsare, D. K. and D. Roy Chowdhuri. Phospholipid distribution in blood and tissues of some submammalian species, 21
 Benjamin, Ronna. See Nevin Weaver
 Bernstein, H. J. See K. Schaumburg
 Berry, James F. and Beverly Kaye. Short communication re chromatography of lipids on polystyrene gel columns, 356
 Bickerstaffe, Roy and James F. Mead. Metabolism of chimyl alcohol and phosphatidyl ethanolamine in the rat brain, 317
 Bierenbaum, M. L. See A. I. Fleischman
 Boch, R. See Marius Lepage
 Brandt, A. E. and W. E. M. Lands. Quantitative gas chromatography, using retention times, 178
 Breckenridge, W. C. and A. Kuksis. Structure of bovine milk fat triglycerides. I. Short and medium chain lengths, 291
 Brenner, R. R. See E. M. Romero
 Brockerhoff, H., R. J. Hoyle, P. C. Hwang and Carter Litchfield. Positional distribution of fatty acids in depot triglycerides of aquatic animals, 24
 Brodnitz Michael H., Wassef W. Nawar and Irving S. Fagerson. Autoxidation of saturated fatty acids. I. The initial products of autoxidation of methyl palmitate, 59
 — Wassef W. Nawar and Irving S. Fagerson. Autoxidation of saturated fatty acids. II. The determination of the site of hydroperoxide groups in autoxidizing methyl palmitate, 65
 Brooks, C. J. W., E. C. Horning and J. S. Young. Characterization of sterols by gas chromatography-mass spectrometry of the trimethylsilyl ethers, 391
 Brown, Ann B. See D. L. Turner
 Butkus, Antanas. See L. Allen Ehrhart

C

- Capella, P., C. Galli and R. Fumagalli. Hydroxy fatty acids from cerebrospines of the central nervous system: GLC determination and mass spectrometric identification, 431
 Carls, Glen A. See Michael M. Martin
 Carr, Sheila. See A. L. Prensky

- Carrion, Maria. See Robert J. Morin
 Chan, Tak. See Giuma M. Sheriha
 Chapman, L. R. See D. F. Kuemmel
 Chicoye, E., W. D. Powrie and O. Fennema. Isolation and characterization of cholesterol-5 β ,6 β -oxidate from an aerated aqueous dispersion of cholesterol, 335
 — W. D. Powrie and O. Fennema. Synthesis, purification and characterization of 7-ketocholesterol and epimeric 7-hydroxycholesterols, 551
 Ching, Te May. Intracellular distribution of lipolytic activity in the female gametophyte of germinating Douglas fir seeds, 482
 Chowdhuri, D. Roy. See D. K. Belsare
 Chung, Li Hsin and Jacqueline Dupont. Acetoacetate metabolism of rats fed high fat or restricted calorie diets, 545
 Cohen, Louis and Juliana Djordjevich. Human serum α -1-lipoprotein patterns revealed by starch gel electrophoresis, 420
 Cornatzer, W. E. See Jerry Baldwin
 Cornwell, David G. See S. Ramachandran
 — See P. Venkata Rao
 Couch, J. R. See A. U. Alam
 Creger, C. R. See A. U. Alam

D

- Dain, Joel A. See John A. Yiamouyiannis
 Delbruck, M. and G. Meissner. Letter to the editor re a note on the availability of phycomyces mutants deficient in β -carotene biosynthesis, 558
 DeLeon, Stella. See N. Nicolaides
 Dillard, C. J. See A. L. Tappel
 Dils, R. See Rodney Watts
 Djordjevich, Juliana. See Louis Cohen
 Dobiasova, Milada and Norman S. Radin. Uptake of cerebroside, cholesterol and lecithin by brain myelin and mitochondria, 439
 Downing, Donald T. and Richard S. Greene. Rapid determination of double-bond positions in monoenoic fatty acids by periodate-permanganate oxidation, 96
 Dryer, R. L. See John R. Paulsrud
 Dupont, Jacqueline. See Li Hsin Chung

E

- Earle, P. R. See Roger W. Miller
 Eggen, Douglas A. See Irwin L. Shapiro
 Ehrhart, L. Allen, Antanas Butkus, A. Lazzarini Robertson, Jr., and Irvine H. Page. Effects of experimental endogenous hyperlipemia on circulating leukocytes and erythrocytes, 34
 Eigen, E. See C. K. Farekh
 Ellingson, John S. and William E. M. Lands. Phospholipid reactivation of plasmalogen metabolism, 111
 Eng, Lawrence F. and Ernest P. Noble. The maturation of rat brain myelin, 157

F

- Fagerson, Irving S. See Michael H. Brodnitz
 Fennema, O. See E. Chicoye
 Fisher, Mary P. and Lawrence A. Wishner. Autoxidation of tissue lipids. II. Monocarbonyl compounds formed by the autoxidation of methyl eicosapentaenoate, methyl docosahexaenoate, and cod-liver oil, 88

- Fleischman, Alan I., Thomas Hayton, Marvin L. Bierenbaum and Portia Watson. The effect of a polyunsaturated diet upon adipose-tissue fatty acids in young coronary males. A five-year cohort study 147
 — H. Yacowitz, M. L. Bierenbaum and T. Hayton. Strain differences in the hypolipemic action of dietary calcium in mature male rats, 1
 Fredrickson, D. S. See A. M. Gotto
 Freidinger, R. M. See K. L. Mikolajczak
 Fumagalli, R. See P. Capella

G

- Galli, C. See P. Capella
 Gans, Joseph H. The distribution of ^{14}C -labeled cholesterol in the dog: Effect of long-term epinephrine administration, 324
 Garton, G. A. See A. K. Lough
 Gold, Martin. Triglyceride sub-classes of various dog adipose tissue sites, 539
 Gotto, A. M., R. I. Levy and D. S. Fredrickson. Preparation and properties of an apoprotein derivative of human serum β -lipoprotein, 463
 Greene, Richard S. See Donald T. Downing

H

- Haigh, W. G., L. J. Morris and A. T. James. Acetylenic acid biosynthesis in *Crepis rubra*, 307
 Hammond, E. W. See L. J. Morris
 Harkness, S. H. See R. Cecil Jack
 Harris, Jeanne. See W. H. Tallent
 Hayton, T. See A. I. Fleischman
 Heath, Robert L. See Kenneth D. Lunan
 Hedin, P. A. See A. C. Thompson
 Hedson, R. D. See A. C. Thompson
 Hempel, Judith E. See Leland L. Smith
 Henry, John E. See Charles L. Sloan
 Henson, R. D. See A. C. Thompson
 Hill, Edward E., William E. M. Lands and Sister P. M. Slakey. The incorporation of ^{14}C -glycerol into different species of diglycerides and triglycerides in rat liver slices, 411
 Hof, Liselotte. See Norman S. Radin
 Holburn, R. R. See D. L. Turner
 Horning, E. C. See C. J. W. Brooks
 Horrocks, Lloyd A. See Grace Y. Sun
 Howse, D. C. and J. R. Wherrett. Short communication re occurrence of inosine in unswathed lipid extracts, 184
 Hoyle, R. J. See H. Brockerhoff
 Hutchins, Roderick F. N. and Michael M. Martin. The lipids of the common house cricket, *Acheta domestica* L. I. Lipid classes and fatty acid distribution, 247
 — The lipids of the common house cricket, *Acheta domestica* L. II. Hydrocarbons, 250
 Hwang, P. C. See H. Brockerhoff

J

- Jack, R. Cecil and S. H. Harkness. Relative incorporation of acetate and glucose into glycerides of *Glomerella cingulata*, 211
 — and Joan A. Laredo. Short communication re fungal spore phospholipids and the accumulation of selected chemicals, 459
 Jackson, Larry L. See Charles L. Sloan
 James, A. T. See W. G. Haigh
 Jantremaky, Judith A. See Irwin L. Shapiro
 Jensen, R. G. See T. A. Marks
 — See C. E. Olney
 — See J. Sampugna
 Johnston, Norah C. See Nevin Weaver

K

- Kampine, John P., Eric Martensson, Ronald A. Yankee and Julian N. Kanfer. Sphingolipid metabolism in leucocytes. I. Incorporation of ^{14}C -glucose and ^{14}C -galactose into glycosphingolipids by intact human leucocytes, 151
 Kanfer, Julian N. See John P. Kampine

LIPIDS, VOL. 3, NO. 6

- Karlsson, K.-A., K. Nilsson and I. Pascher. Separation of lipids containing phytanic acid by thin-layer chromatography, 389
 Kaye, Beverly. See James F. Berry
 Kircher, Henry W. Short communication re an inexpensive, unbreakable chromatographic column, 105
 Kleiman, R. See C. R. Smith, Jr.
 Klopffenstein, W. E. and Y. Pomeranz. Fatty acids in lipids of maturing wheat, 557
 Knipprrath, Werner G. and James F. Mead. The effect of the environmental temperature on the fatty acid composition and on the *in vivo* incorporation of ^{14}C -acetate in goldfish (*Carassius auratus* L.), 121
 Knoche, H. W. A study on the biosynthesis of *cis*-9,10-epoxyoctadecanoic acid, 163
 Knudson, Alfred G., Jr. See George Rouser
 Kohler, Daniel P. See Leland L. Smith
 Kritchevsky, David. See Irwin L. Shapiro
 — Shirley A. Tepper and George H. Rothblat. Short communication re the effect of bile salts on hydrolysis of cholesterol oleate by rabbit aorta, 454
 Kritchevsky, Gene. See George Rouser
 Kuemmel, D. F. and L. R. Chapman. The 9-hexadecanoic and 11-octadecanoic acid content of natural fats and oils, 313
 Kuksis, A. See W. C. Breckenridge

L

- Lambreton, E. N. and Randall Wood. Glyceryl ethers in insects: Identification of alkyl and alk-1-enyl glyceryl ether phospholipids, 503
 Lands, William E. M. See A. E. Brandt
 — See John S. Ellingson
 — See Edward E. Hill
 — See Sister P. M. Slakey
 Laredo, Joan A. See R. Cecil Jack
 Law, John H. See Nevin Weaver
 Lepage, Marius. The lipid components of white potato tubers (*Solanum tuberosum*), 477
 — and R. Koch. Pollen lipids attractive to honey bees, 539
 Levy, R. I. See A. M. Gotto
 Libbey, L. M. and J. P. Walradt. Short communication re 3,5-di-*tert*-4-hydroxytoluene (BHT) as an artifact from diethyl ether, 557
 Lim, James. See Edward O. Oswald
 Litchfield, Carter. Predicting the positional distribution of docosahexaenoic and docosapentaenoic acids in aquatic animal triglycerides, 417
 — Triglyceride analysis by consecutive liquid-liquid partition and gas-liquid chromatography. *Ephedra nevadensis* seed fat, 170
 — See H. Brockerhoff
 Loeblich, Alfred R., III, and V. Elliott Smith. Chloroplast pigments of the marine diflagellate *Gyrodinium aureolum*, 5
 Lough, A. K. and G. A. Garton. The lipids of human pancreas with special reference to the presence of fatty acid methyl esters, 321
 Lowrey, Robert R. and Ian J. Tinsley. Short communication re a device to apply detection reactions to microplates, 106
 Lunan, Kenneth D. and Robert L. Heath. Short communication re irreversible enzyme inhibition by a phosphatidic acid-like lipid, 456
 Lyman, Richard L. See Rosemarie Ostwald

M

- MacDonald, Robert C. and James F. Mead. The α -oxidation system of brain microsomes. Cofactors for α -hydroxy acid decarboxylation, 275
 Marks, T. A., J. G. Quinn, J. Sampugna and R. G. Jensen. Studies on the specificity of a lipase system from *Geotrichum candidum*, 143
 Marshall, M. O. See L. J. Morris
 Martensson, Eric. See John P. Kampine
 Martin, Michael M. See Roderick F. N. Hutchins
 — and Glen A. Carls. The lipids of the common house cricket, *Acheta domestica* L. III. Sterols, 255
 Matthews, W. Stephen and Leland L. Smith. Sterol metabolism. III. Sterols of marine waters, 239
 Mead, James F. See Roy Bickerstaffe
 — See Werner G. Knipprrath
 — See Robert C. MacDonald
 Medwadowski, B. F. See J. Van der Veen
 Meisner, G. See M. Delbruck
 Mikolajczak, K. L., R. M. Freidinger, C. R. Smith, Jr., and I. A. Wolff. Oxygenated fatty acids of oil from sunflower seeds after prolonged storage, 489
 — C. R. Smith, Jr. and I. A. Wolff. Glyceride structure of *Cardamine hirsuta* L. seed oil, 215

Miller, Roger W., F. R. Earle, I. A. Wolff and A. S. Barclay. Search for new seed oils. XV. Oils of Boraginaceae, 43

Minyard, J. P. See A. C. Thompson

Morin, Robert J. and Maria Carrion. In vitro incorporation of acetate-1-¹⁴C into the phospholipids of rabbit and human endometria, 349

Morris, L. J. Fatty acid composition of *Claviceps* species. Occurrence of (+)-three- β , 10-dihydroxystearic acid, 260

— See W. G. Haigh

— M. O. Marshall and E. W. Hammond. The trans-3-enoic acids of *Aster alpinus* and *Arctium minus* seed oils, 91

Morrison, W. R. The distribution of phospholipids in some mammalian milk, 101

— Fatty acid composition of milk phospholipids. III. Camel, ass and pig milks, 107

N

Nair, P. P. See N. Nicolaides

Nawar, Wasef W. See Michael H. Brodnitz

Nelson, Gary J. Short communication re differentiation of nitrogenous phospholipids by infrared absorption between 9 and 11 microns, 104

— Studies on the lipids of sheep red blood cells. II. The incorporation of phosphorus into phospholipids of HK and LK cells, 267

Nichols, B. W. Fatty acid metabolism in the chloroplast lipids of green and blue-green algae, 354

— and B. J. B. Wood. The occurrence and biosynthesis of gamma-linoleic acid in a blue-green alga, *Spirulina platensis*, 46

Nicolaides, N. and M. N. A. Ansari. Fatty acids of unusual double-bond positions and chain lengths found in rat skin surface lipids, 403

— P. P. Nair and Stella DeLeon. Short communication re the occurrence of 5 α -cholestan-3 β -ol (dihydrocholesterol) in human skin surface lipid, 458

Nilsson, K. See K.-A. Karlsson

Nishimura, Kenji and Tamio Yamakawa. Isolation of cerebroside containing glucose (glucosyl ceramide) and its possible significance in ganglioside synthesis, 262

Noble, Ernest P. See Lawrence F. Eng

O

Olcott, H. S. See J. Van der Veen

— and J. Van der Veen. Comparison of antioxidant activities of tocopherol and its methyl derivatives, 331

Olney, C. E., R. G. Jensen, J. Sampugna and J. G. Quinn. The purification and specificity of a lipase from *Veronica anthelmintica* seed, 498

Ostwald, Rosemarie and Richard L. Lyman. Influence of sex and gonadal hormones on lipid metabolism in essential fatty acid-deficient rats, 199

Oswald, Edward O., Carl E. Anderson, Claude Piantadosi and James Lim. Metabolism of alkyl glyceryl ethers in the rat, 51

P

Page, Irvine H. See L. Allen Ehrhart

Parekh, C. K. and E. Eigen. Tritium and ¹⁴C counting in tissue samples by using lipid scintillation method, 225

Paecher, I. See K.-A. Karlsson

Paulsrud, John R. and R. L. Dryer. Circum-annual changes in triglyceride fatty acids of bat brown adipose tissue, 340

Pawar, Sitaram S. Short communication re in vitro effect of prostaglandin (PGE₁) on the release of glycerol and the metabolism of palmitic acid in rat adipose tissue, 383

— and Herbert C. Tidwell. Dietary fat effect on incorporation and release of lipids and cholesterol by rat intestinal slices, 346

Pelick, Nicholas. See Walter R. Supina

Perkins, E. G. See C. J. Argoudelis

Phillips, Gerald B. Short communication re fatty acids of human red cell total phospholipids, 385

Piantadosi, Claude. See Edward O. Oswald

Pomeranz, Y. See W. E. Klopfenstein

Powrie, W. D. See E. Chicoye

Prensky, A. L. and Sheila Carr. Short communication re the behavior of proteolipids on dextran gel columns eluted with organic solvents, 453

Q

Quinn, J. G. See T. A. Marks

— See C. E. Olney

R

Radin, Norman S. See Milada Dobiasová

— Liselotte Hof and Carolyn Seidl. Short communication re lipid contaminants: Polypropylene apparatus and vacuum pumps, 192

Ramachandran, S. See P. Venkata Rao

— Howard W. Sprecher and David G. Cornwell. Studies on the preparation and analysis of glyceryl ether derivatives and the isolation and reductive ozonolysis of unsaturated glyceryl ethers, 511

Rao, P. Venkata, S. Ramachandran and David G. Cornwell. Short communication re thin-layer chromatography of phospholipids on alumina, 187

Renkonen, Ossi. Short communication re individual molecular species of phospholipids. VII. Analysis of lecithins containing ten to twelve double bonds, 191

Robertson, A. Lazzarini, Jr. See L. Allen Ehrhart

Romero, E. M. and R. R. Brenner. Short communication re the phosphatidyl ethanolamine of *Pseudomonas aeruginosa* grown in hexadecane, 460

Rose, Lewis P. See Walter R. Supina

Rothblat, George H. See David Kritchevsky

Rouser, George, Gene Kritchevsky, Akira Yamamoto, Alfred G. Knudson, Jr. and Gerald Simon. Short communication re accumulation of a glycerolphospholipid in classical Niemann-Pick disease, 287

— and Akira Yamamoto. Short communication re curvilinear regression course of human brain lipid composition changes with age, 284

S

Sampugna, J. See T. A. Marks

— See C. E. Olney

— and R. G. Jensen. Suitability of *Geotrichum candidum* lipase for the stereospecific analysis of some triglycerides, 519

Schaumburg, K. and H. J. Bernstein. Calculation of the NMR spectrum of double-bond protons in aliphatic systems, 193

Seidl, Carolyn. See Norman S. Radin

Shapiro, Irvin L., Judith A. Jastremsky, Douglas A. Eggen and David Kritchevsky. Cholesterol metabolism in the baboon, 135

— Judith A. Jastremsky and David Kritchevsky. Short communication re effect of hypercholesterolemia on the activity of serum lecithin-cholesterol acyltransferase, 381

Sheriha, Gluma M., George R. Waller, Tak Chan and Allen D. Tillman. Composition of bile acids in ruminants, 72

Sliver, M. J. See D. L. Turner

Simon, Gerald. See George Rouser

Slakey, Sister F. M. See Edward E. Hill

— and W. E. M. Lands. The structure of rat liver triglycerides, 30

Sloan, Charles L., Larry L. Jackson, Graeme L. Baker and John E. Henry. Short communication re fatty acid methyl esters in grasshopper eggs, 455

Smith, C. R., Jr. See K. L. Mikolajczak

— R. Kleiman and I. A. Wolff. *Caltha palustris* L. seed oil. A source of four fatty acids with cis-5-unsaturation, 37

Smith, Leland L. See W. Stephen Matthews

— Daniel P. Kohler, Judith E. Hempel and Johan E. van Lier. Sterol metabolism. IV. Microbial disposition of 5 β cholesterol-3 β -ol, 301

Smith, V. Elliott. See Alfred R. Loeblich, III

Snyder, Fred. See Randall Wood

Spencer, G. F. See W. H. Tallent

Sprecher, Howard W. The total synthesis and metabolism of 4-decenoate dodeca-3,6-dienoate, tetradeca-5,8-dienoate and hexadeca-7,10-dienoate in the fat-deficient rat, 14

— See S. Ramachandran

Sun, Grace Y. and Lloyd A. Horrocks. The fatty acid and aldehyde composition of the major phospholipids of mouse brain, 79

Supina, Walter R., Nicholas Pelick, Lewis P. Rose and Gary C. Walker. Short communication re selection and evaluation of gas chromatographic stationary phases for qualitative separation of components of lipid mixtures, 374

T

- Tallent, W. H., Jeanne Harris, G. F. Spencer and I. A. Wolff. Structure and intraglyceride distribution of coriolic acid, 425
- Tamaki, Yoshio. Short communication re isolation of tetradecan-1,14-dioic acid from the Comstock mealybug, *Pseudococcus comstocki* Kuwana (Homoptera: Pseudococcidae), 186
- Tappel, A. L. and C. J. Dillard. Retinol inhibition of some proteolytic enzymes, 221
- Tepper, Shirley A. See David Kritchevsky
- Thompson, A. C., R. D. Henson, P. A. Hedin and J. P. Minyard. Constituents of the cotton bud. XII. The carotenoids in the buds, seeds and other tissue, 495
- R. D. Henson, J. P. Minyard and P. A. Hedin. Short communication re fatty acid composition of polar lipids of cotton buds, 373
- Tidwell, Herbert C. See Sitaram S. Pawar
- Tillman, Allen D. See Giama M. Sheriha
- Turner, D. L., M. J. Silver, R. R. Holburn and E. Baczynski. The total synthesis of phosphatidyl(dioleoyl)hydroxy-L-proline and its activity in blood-clotting systems, 228
- M. J. Silver, R. R. Holburn, E. Baczynski and Ann B. Brown. A unsaturated phosphonic acid analogue of phosphatidylethanolamine and its activity in blood-clotting systems, 234
- Tyrrell, David. The fatty acid composition of some entomophthoraceae. II. The occurrence of branched-chain fatty acids in *Conidiobolus denaesporus* Drechs., 368

V

- Van der Veen, J. See H. S. Olecott
- B. F. Medwadowski and H. S. Olecott. Short communication re losses of fatty acids during the saponification extraction of small samples, 189
- van Golde, L. M. G. See Moseley Waite
- van Lier, Johan E. See Leland L. Smith

W

- Waite, Moseley and L. M. G. van Golde. Dietary induced alterations in swelling characteristics and endogenous phospholipase A₂ activity of rat liver mitochondria, 449
- Walker, Gary C. See Walter R. Supina
- Waller, George R. See Giama M. Sheriha
- Walratt, J. P. See L. M. Libbey
- Ware, Camellia M. and Lawrence A. Wishner. Short communication re the lipid antioxidant properties of iodine compounds, 182
- Watson, Portia. See Alan I. Fleischman
- Watts, Rodney and R. Dils. Human milk: Quantitative gas-liquid chromatographic analysis of triglyceride and cholesterol content during lactation, 471
- Weaver, Nevin, Norah C. Johnston, Ronna Benjamin and John H. Law. Novel fatty acids from the royal jelly of honeybees (*Apis mellifera*, L.), 535
- Wherrett, J. R. See D. C. Howse
- Wishner, Lawrence A. See Mary P. Fisher
- See Camellia M. Ware
- Wolff, I. A. See K. L. Mikolajczak
- See Roger W. Miller
- See C. R. Smith, Jr.
- See W. H. Tallent
- Wood, B. J. B. See B. W. Nichols
- Wood, Randall. See E. N. Lambremont
- and Fred Snyder. Quantitative determination of alk-1-enyl- and alkyl-glycerol ethers in neutral lipids and phospholipids, 129

X-Y-Z

- Yacowitz, H. See A. I. Fleischman
- Yamakawa, Tamio. See Kenji Nishimura
- Yamamoto, Akira. See George Rouser
- Yankee, Ronald A. See John P. Kampine
- Yiamouyiannis, John A. and Joel A. Dain. Short communication re UDP-galactose: N-acetylgalactosaminyl-(N-acetylneuraminyl) galactosyl-glucosyl-ceramide transferase activity in adult frog brain, 378
- Young, J. S. See C. J. W. Brooks



SUBJECT

A

- Abetalipoproteinemia, fatty acids in erythrocytes, 387
- Bis(trimethylsilyl)acetamide, in silylation of triglyceride hydrolyzates for GLC, 425
- Acetate, incorporation into fatty acids of fish, 121
 - incorporation into fungal glycerides, 211
 - utilization after fat diets, rat intestinal slices, 346
- Acetic acid, incorporation into fatty acids of *Crepis rubra*, 307
- 1-¹⁴C-Acetic acid, incorporation and oxidation by rabbit endometrium, 349
- Acetoacetic acid, incorporation into cholesterol, rat, 545
 - incorporation in fatty acids, rat, 545
 - oxidation by rat, 545
 - rat, dietary fat effect on metabolism, 545
- Acetoxyhydroxy acid, methyl ester, IR, NMR, mass, 215
- 13(14)-Acetoxy-14(13)-hydroxydocosanoic acid, *Cardamine impatiens* seed oil, 215
 - mass spectrum, 215
 - synthesis, 215
- 15(16)-Acetoxy-16(15)-tetracosanoic acid, *Cardamine impatiens* seed oil, 215
- Acetylated dihydroxy acids, *Cardamine impatiens* triglycerides, 215
- β -Acetylglucosaminidase EC 3.2.1.30, effect of lipids, 221
- Acetylenic fatty acid, biosynthesis by *Crepis rubra*, 307
- Acheta domestica, fatty acid profile, 247
 - hydrocarbons in cricket, 250
 - lipid classes, 247
 - sterols, 256
- ACP. See acyl carrier protein
- Activated sludge, treatment of sewage, sterol removal, 301
- N-Acyl neuraminic acid. See also sialic acid
- Adrenal, dog, cholesterol distribution between tissue and plasma, 324
- Adenosine triphosphate, effect on α -oxidation, 275
- Adipose tissue, brown, of bat, 340
- Adipose tissue, human, fatty acid profile, 147
 - fatty acids, effect of polyunsaturated diet, 147
 - fatty acids, GLC of, 147
 - fatty acids, linoleic acid concentration, 147
 - fatty acids, stable pool, 147
 - fatty acids, turnover time, 147
- Adipose tissue metabolism, rat, PGE, effect, 383
- Adsorption of lipids, effect of salts, 439
- Aerated cholesterol dispersion, oxidation products, 335
- Age, changes in human brain lipid with, 284
 - effect on incorporation of precursors into fungal glycerides, 211
- Albumin, staining with oil red O, 420
- Aldehydes, profile of mouse brain phospholipids, 79
 - See also fatty aldehydes; plasmalogens
- Algae, *Anabaena cylindrica* blue-green algae, fatty acid metabolism in, 354
 - *Anacystis nidulans*, blue-green algae, fatty acid metabolism in, 354
 - *Chlorella vulgaris* green algae, fatty acid metabolism in, 354
 - galactosyl diglycerides, 354
 - growth media, 354
 - phosphatidyl glycerol, 354
 - photosynthesis, 354
- Algae. See microorganisms
- Alkanes, in crickets, 250
- Alkenyl acyl ethanamine phosphoglycerides, biosynthesis, rats, 317
- Alkenyl ether, structure relation to imidazole, 111
 - TLC, 129
- Alkenyl glycerol-3-phosphorylcholine, alkenyl hydrolase, 111
- Alkenyl hydrolase, 111
- Alk-1-enyl ethers, TLC, 129
- Alk-1-enyl glyceryl ethers, occurrence in insects, 503
- Alkyl acyl ethanamine phosphoglycerides, biosynthesis, rats, 317
- Alkyl ethers, TLC, 129
- Alkyl glyceryl ethers, occurrence in insects, 503
 - rat metabolism of, 51
- Alkyl glycerols, profile, location of olefinic bonds, 511
 - See glycerol ethers
- Alkyl monoolides and diolides, glyceryl ethers, ozonolysis products, 511
- Alkyl (C⁹ to C¹⁹) tolyl methyl trimethyl ammonium chloride. See hyamine
- Alternaria oleracea, fungal spore phospholipids, fatty acid profile, 459
- Alumina, TLC, phospholipids, 187
- Amaretic familial idiocy, late infantile, 287
 - See gangliosidosis

- American cockroach. See insect
- American eel, triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
- Amphibia. See specific name
- Amphibian, triglycerides, positional distribution of fatty acids, 24
- Amsinckia tessellata Gray, fatty acid profile, properties, 48
- Anabaena cylindrica, fatty acid profiles, total lipids and neutral lipids, 46
 - See algae
- Anabaena flos-aquae, fatty acid profiles, total lipids and neutral lipids, 46
- Anacystis nidulans, fatty acid profiles, total lipids and neutral lipids, 46
 - See algae
- Anthonomus grandis Boheman. See insect
- Ancistrodon piscivorus piscivorus, phospholipase activity, 107
- Anchusa angustifolia L., fatty acid profile, properties, 43
- Anchusa azurea Mill., fatty acid profile, properties, 43
- Anchusa leptophylla Roem. & Schult., fatty acid profile, properties, 43
- Anchusa officinalis L., fatty acid profile, properties, 43
- Anchusa, fatty acid profile, properties 43
- Androsterone, GLC, 374
- Annual change, in lipid comp of human brain, 284
- Antheraxanthin, of Euglena gracilis, 5
 - possible identity with diadinoxanthin, 5
- Anthonomus grandis. See insects
- Anticoagulant activity, of phosphatidyl hydroxy-L-proline, 228
- Antioxidant, property of iodine compounds, 182
- Antioxidant activity, 7,8, and 5,7,8-trimethyl and γ -methyl tocopherol, 331
- Antioxidant deficiency. See also α -tocopherol deficiency
- Aorta, dog, cholesterol distribution between tissue and plasma, 324
 - rabbit, hydrolysis of cholesterol oleate, 454
- Apis mellifera L. See insects
- Apoprotein, from human serum β -lipoprotein, solubilization by succinylation, 463
- Aquatic animals, triglycerides, positional distribution of fatty acids, 24
 - triglyceride species, 417
- Arachidonic acid, rabbit liver microsomal phospholipids, variation with age, 361
- Arachidonic acid (20:4 ω 6), biosynthesis in rats, 199
- Arctium minus. See oilseeds
- Arnebia griffithii Boiss., fatty acid profile, properties, 43
- Arylsulfatase EC 3.1.6.1, effect of lipids, 221
- Asclepias syriaca, isomer distribution in monoenoic esters, 313
- Ascorbic acid, cofactor for α -oxidation, 275
- Ass milk, phospholipids, fatty acid profiles, 107
 - phospholipids profile, 101
- Aster alpinus. See oilseeds
- ATP. See adenosine triphosphate
- Auroxanthin, from cotton plants, properties, 495
- Automated analysis, enzymes inhibited by lipids, 221
- Autooxidation, of cod liver oil, 88
 - of palmitic acid, methyl ester, 59
 - of tissue lipids, 88
- Autoxidation, of oils, squalene, 331
- Autoxidation of sterols, 239
- Avocado, isomer distribution in monoenoic esters, 313

B

- Baboon, cholesterol metabolism, 136
- Barndoor skate, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
- Rat brown adipose tissue, fatty acids, triglycerides, 340
- α -Batyl alcohol (1-octadecyl glyceryl ether), rat metabolism of, conversion to ether phospholipids, 51
- β -Batyl alcohol (2-octadecyl glyceryl ether), rat metabolism of, conversion to ether phospholipids, 51
- Beef, isomer distribution in monoenoic esters, 313
 - See bovine
- Bile, dog, cholesterol distribution between tissue and plasma, 324
- Bile acids, comp in fetus bile, ruminants, 72
 - comp in ruminants, 72
- Bile salts, effect on aortic hydrolysis of cholesterol oleate, 454
- Biosynthesis, liver, diglycerides and triglycerides, 411

- Birds.** See specific name
 — triglycerides, positional distribution of fatty acids, 24
Blood, red cells. See erythrocyte
Blood coagulation, effect of dioleoylglycerol(2-amino-ethyl)phosphonate, 234
 — effect of phosphatidyl ethanolamine (as dioleoyl and diinoleoyl compounds), 234
 — effect of phosphatidyl hydroxy-L-proline on, 228
Blood lipid and cholesterol levels, effect of fat diets, rat, 346
Blood serum, in Triton hyperlipemia, dog, 84
Boll weevil. See insects
Boraginaceae, fatty acid profile, properties, 43
Bovine, bile acids, comp, 72
 — feces, bile acids, comp, 72
Bovine milk, phospholipids profile, 101
Bovine milk, triglyceride profile, 291
Brain cerebroside, hydroxy fatty acids, 431
Brain, fish, (*Heteropneustes fossilis*), phospholipids, 21
 — frog, ganglioside biosynthesis, *in vitro*, 378
 — human, annual change in comp, 284
 — human, lipid comp, age changes, 284
 — pigeon, (*Columba livia*), phospholipids, 21
 — mouse, phospholipids, fatty acid profile, 79
 — mouse, phospholipids, fatty aldehyde profile, 79
 — rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
 — rat, lipid adsorption by particle, 439
 — rat, metabolism, chimyl alcohol, 317
 — rat, metabolism, phosphatidyl ethanolamine, 317
 — rat, myelin development, 157
 — rat, α -oxidation system in microsomes, 275
 — toad, (*Bufo malanostictus*), phospholipids, 21
 — turtle, (*Kachuga Smithi*), phospholipids, 21
Bromelain EC-3.4.4.24, effect of lipids, 221
1-Bromo-2-octyne, intermediate in unsaturated fatty acid synthesis, 14
Brown adipose tissue, bat, triglyceride comp, 340
Buffers, for lipoprotein electrophoresis, 420
Bufo malanostictus. See toad
Burbot (maria), triglycerides, positional distribution of fatty acids, 24
3,5-di-tert-Butyl-4-hydroxytoluene, as contaminant from diethyl ether, 557

C

- Calcium,** dietary, effect on rat hypolipemia, 1
Caltha palustris L. See oilseed
Camel, isomer distribution in monoenoic esters, 313
Camel milk, phospholipids, fatty acid profiles, 107
 — phospholipids profile, 101
A5,7-Campestadiene-3 β -ol, in crickets, 256
Campestene-3 β -ol, in crickets, 256
A5-Campestene-3 β -ol, in crickets, 256
A7-Campestene-3 β -ol, in crickets, 256
Campesterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 — in marine waters, GLC TLC, 239
Carassius auratus L. See goldfish
Carbon-14, measurement in tissues, 225
Carbon dioxide from fatty acids, FGE effect on rat adipose tissue, 383
Carbonyl reduction, 5
Cardamine impatiens seed oil, triglycerides, 215
Cardiolipin, human brain, equations for computation, 284
 — TLC, 187
 — See also diphosphatidyl glycerol
Carnitin, identification as inosine, 184
Carotenes, inhibition of enzymes, 221
 — partition coefficients, 5
 α -Carotene, from cotton plants, properties, 495
 β -Carotene, of *Gyrodinium resplendens*, 5
 — spectra, ultra-violet, visible; column chromatography, 5
 β -Carotene mutants, of *Phycomyces*, 558
Carotenoids, from cotton bud, leaf, petal, seed, seedling, 495
 — of *Gyrodinium resplendens*, 5
 — pollen lipids attractive to honey bees, 530
 — in potato tubers, 477
Castor bean lipase, hydrolysis of triglycerides containing dihydroxy acyl groups, 215
Cathepsin-D EC-3.4.4.23, effect of lipids, 221
CCD. See countercurrent distribution
Ceramides, N-acyl sphingosine. See also sphingosine cr under glycoside name
 — human brain, equations for computation, 284
Cerebroside, adsorption by brain particles, 439
 — human brain, equations for computation, 284
 — hydroxy fatty acids, 431
 — from potato tubers, 477
Chicken serum, phosphatidyl choline-cholesterol acyl-transferase, 381
Chimyl alcohol, metabolism in rat brain, 317
 α -Chimyl alcohol (1-hexadecyl glyceryl ether), rat metabolism of, conversion to ether phospholipids, 51
 β -Chimyl alcohol (2-hexadecyl glyceryl ether), rat metabolism of, conversion to ether phospholipids, 51
Chlorella vulgaris. See algae
Chloroglossa fritschii, fatty acid profiles, total lipids and neutral lipids, 46
Chlorophyll a, of *Gyrodinium resplendens*, 5
 — spectra, ultra-violet, visible, 5
 — spectra, ultra-violet, visible; column chromatography, 5
Chlorophyll c, of *Gyrodinium resplendens*, 5
 — spectra, ultra-violet, visible, 5
 — spectra, ultra-violet, visible; column chromatography, 5
Chloroplast lipids, fatty acid metabolism in, 354
 — metabolism in green and blue-green algae, 354
Chloroplast pigment Gy 439, spectra, ultra-violet, visible; column chromatography, 5
Chloroplast pigment Gy 442, spectra, ultra-violet, visible; column chromatography, 5
Chloroplast pigment Gy 443 (a + b), spectra, ultra-violet, visible; column chromatography, 5
Chloroplast pigments, of marine dinoflagellate (*Gyrodinium resplendens*), 5
3-Keto-5 β -Cholanoic acid, in bile, sheep, bovine, fetus; GLC, 72
A5,7-Cholestadiene-3 β -ol, in crickets, 256
A5,22-Cholestadiene-3 β -ol, in crickets, 256
5 α -Cholestan-2 β -ol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
5 α -Cholestan-3 α -ol, dihydrocholesterol, 458
 — GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
5 α -Cholestan-3 β -ol, GLC conditions and derivatives for GLC, 458
 — GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 — in human skin surface lipids, 458
5 β -Cholestan-2 β -ol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
5 β -Cholestan-3 β -ol, from domestic sewage, 391
 — GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Cholestane-3 β -ol, in crickets, 256
A5,7,22-Cholestatriene-3 β -ol, in crickets, 256
Cholest-5-ene-3 β ,24 α -diol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Cholest-5-ene-3 β ,25-diol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
A5-Cholestene-3 β -ol, in crickets, 256
A7-Cholestene-3 β -ol, in crickets, 256
Cholest-4-en-3 β -ol, GLC retention times for trimethylsilyl ether, principal ions in mass spectra, 391
5 α -Cholest-7-en-3 β -ol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
5 α -Cholest-8(14)-en-3 β -ol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Cholest-4-en-3 β -yl trimethylsilyl ethers, mass spectrum, 391
5 α -Cholest-7-en-3 β -yl trimethylsilyl ethers, mass spectrum, 391
5 α -Cholest-8(14)-en-3 β -yl trimethylsilyl ethers, mass spectrum, 391
Cholest-5-en-3 β -ol-7-one, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Cholest-5-en-3 β -ol-24-one, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Cholesterol, adsorption by brain particles, 439
 — biosynthetic rate and turnover time in the baboon, 136
 — complex with phospholipids, with galactolipid, 157
 — from domestic sewage, 301
 — exogenous and biosynthetic, transport by serum alpha- and beta-lipoproteins in the baboon, 136
 — GLC conditions and derivatives for GLC, 458
 — GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 — human brain, equations for computation, 284
 — in human colostrum and milk, 471
 — influence of sex hormones on EFA deficiency in rats, 199
 — inhibition of enzymes, 221
 — in marine waters, GLC, TLC, 239
 — oxidation by aeration, product, 335
 — pool size in baboon, 136
 — rat brain myelin development, 157
Cholesterol-5 β ,6 α -oxide, isolation and characterization from aerated cholesterol, 335

- 4-¹⁴C-Cholesterol, distribution in dogs, time course, 324
 7 α -³H-Cholesterol, incorporation into serum cholesterol esters in the baboon, 136
 Cholesterol biosynthesis, from acetate; from mevalonic acid, in the baboon, 136
 Cholesterol biosynthesis and release, effect of fat diets, rat, 346
 Cholesterol esters, in human pancreas, 321
 — incorporation of cholesterol; mevalonic acid, in the baboon, 136
 — influence of sex hormones on EFA deficiency in rats, 199
 — human brain, equations for computation, 284
 Cholesteryl oleate, hydrolysis by rabbit aorta, 454
 Cholesteryl phytanate, prep. TLC, 389
 Cholesteryl trimethylsilyl ethers, mass spectrum, 391
 Choline phosphotransferase, in liver, variation with age, rabbit, 361
 6-Chromanol, 2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl), lipid antioxidant property, 182
 6-Chromanol, 2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl). See also α -tocopherol.
 Chromatography, column, of brain cerebroside, 431
 — brain glycolipids (Floridil), 262
 — β -carotene, 5
 — carotenoids (magnesium oxide-Celite), 495
 — chlorophyll a, 5
 — chlorophyll c, 5
 — chloroplast pigment Gy 439, 5
 — chloroplast pigment Gy 442, 5
 — chloroplast pigment Gy 443 (a + b), 5
 — chloroplast pigments (silica gel H-Celatom), 5
 — DEAE, rat brain lipids, 317
 — dextran gel, elution of proteolipids, 453
 — diadinoxanthin, 5
 — diadinoxanthin-furanoid-oxide, 5
 — dinoxanthin, 5
 — dinoxanthin-furanoid-oxide, 5
 — fatty acids methyl esters (saturates, monoenes and dienes) (silicic acid-silver nitrate), 403
 — inexpensive unbreakable column, 105
 — of monoenoic methyl esters (Celite-silica gel-silver nitrate), 313
 — nerve lipids (polystyrene gel), 385
 — neutral lipid separation, 121
 — peridinin, 5
 — peridinin-furanoid-oxide, 5
 — polar and non-polar lipids (polystyrene gel), 385
 — polystyrene gel, 385
 — pyrrhoxanthin, 5
 — pyrrhoxanthin-furanoid-oxide, 5
 — silicic acid (total potato lipids), 477
 — TEAE cellulose for phospholipids, 228
 — triglycerides (Floridil), 30
 Chromatography, gas-liquid, alkyl iodides (silica gel G), 511
 — androsterone, 374
 — band-broadening, 178
 — bile acids, 72
 — capillary, *cis*-hexadecenoic and *cis*-octadecenoic acid methyl ester positional isomers, 313
 — 5 α -cholestan-3 β -ol, 458
 — cholesterol, 458
 — cholesterol, 5 β -cholestan-3 β -ol, 301
 — cholesterol content of human colostrum and milk, 471
 — of cholesterol derivatives, retention times, 551
 — cholesterol-5,6-oxides, 335
 — column characterization, 374
 — column selection and evaluation, 374
 — 5 β -coprostan-3 β -ol, 458
 — correction factor for dtm peak area, 178
 — dehydroepiandrosterone, 374
 — dibasic acid, 186
 — diffusion, 178
 — of diglyceride acetates, containing up to twelve bonds, 191
 — estrogen, 374
 — etiocholanolone, 374
 — fatty acid butyl esters, 291
 — fatty acid methyl esters, 374
 — of fatty acid methyl esters, C₁₅-C₄₀, 403
 — fatty acid methyl ester profile, 455
 — fatty acids, 349
 — fatty acids of potato tubers, 477
 — fatty acid profile of human colostrum and milk, 471
 — 11- β -hydroxyetiocholanolone, 374
 — of hydroxy fatty acids, 431
 — 11-ketotiocholanolone, 374
 — 17-keto-steroids, 374
 — lipids qualitative separation, 374
 — long-chain acetylated dihydroxy acid, methyl ester, 215
 — milk fat anal, 291
 — of mono- and dibasic fatty acids methyl esters, 65
 — new silicone phases, 374
 — peak area correction factor, 178
 — pollen lipids attractive to honey bees, 530
 — preparative for methyl palmitate, 59
 — preparative, triglycerides, 291
 — retention times, 178
 — sewage sterols, 301
 — of silylated enzymatic hydrolysates of triglycerides, 425
 — stationary phases evaluation, OV-1, OV-101, OV-3, OV-7, OV-11, OV-17, QF-1, XE-60, DEGS, 374
 — sterols; cholesterol, stigmasterol, β -sitosterol, 289
 — sterol trimethylsilyl ethers, 391
 — fatty acid profile, 539
 — triglyceride profile in human colostrum and milk, 471
 — triglycerides, 170, 291
 — trimethylsilane derivative of cerebroside sugar, 262
 Chromatography, gas-liquid-mass, hydroxy fatty acids as trimethylsilyl ethers, 431
 Chromatography, gas-liquid-mass spectrometer, di-TMS derivatives of hydroxylated monounsaturated fatty acids, 379
 Chromatography, liquid-liquid partition, partition numbers, 170
 — (reverse phase), preparative, 170
 — (reverse phase), triglycerides, 170
 Chromatography, mass spectrometer. See chromatography, gas-liquid-mass spectrometer
 Chromatography, Sephadex, removal of inosine from lipids, 184
 Chromatography, Sephadex column, purification of lipid extracts, 267
 Chromatography, thin-layer, acetylated dihydroxy acid, methyl ester, 215
 — alkyl- and alk-1-enyl-glycerol ethers, 129
 — alkyl monooxides, alkyl diiodides (silica gel G), 511
 — alumina (phospholipids), 187
 — apparatus for applying reagents to microplates, 105
 — of brain cerebroside, 431
 — carotenoids (kieselguhr coated with vegetable oil), 495
 — cholesterol and 5 β -cholestan-3 β -ol, 301
 — cholesterol derivatives (silica gel), 551
 — cholesterol and other lipids, 346
 — cholesterol oxidation products, 335
 — detection procedure (phospholipids), 111
 — diglycerides (silica gel H boric acid) (silica gel-silver nitrate), 238
 — dihydroxy fatty acids (silica gel G-boric acid), 489
 — diimide reduction of unsaturated fatty acid (silica-gel G-silver nitrate), 489
 — 2-dimensional separation of myelin lipids, 157
 — di- and triglycerides (silicic acid, silicic acid-silver nitrate), 411
 — epoxy fatty acids (silica gel G), 489
 — ethanolamine glycerophosphatide plasmalogen, 157
 — fatty acid methyl esters (silica gel G; silica gel G-boric acid), 260
 — glycolipids (borax-impregnated silica gel), 262
 — glycolipids (silica gel G), 262
 — 32p in individual spots dtm, 267
 — lipid components of potato tubers, 477
 — microplates, microscope slides, 105
 — neutral lipids (silica gel G), 51
 — palmitic acid, methyl ester autooxidized, 59
 — partition numbers, 170
 — phospholipids, 111, 349
 — phospholipids (silica gel H), 51
 — phosphorus recovery, 187
 — phytanic acid containing lipids, 389
 — pollen lipids attractive to honey bees, 530
 — polyunsaturated diglyceride acetates, (silica gel-silver nitrate), 191
 — polyunsaturated dimethyl phosphatides (silica gel-silver nitrate), 191
 — preparative (cellulose, for inosine), 184
 — preparative (cholesterol derivatives), 551
 — preparative (silica gel H-magnesium triisilicate: phospholipids), 107
 — preparative for unsaturated fatty acids (silica gel-silver nitrate), 37
 — preparative (triglycerides), 291
 — quantitative, 129
 — quantitative dtm of myelin lipids, 157
 — sewage sterols, 301
 — solvent system (carotenoids and epoxy fatty acids), 163
 — solvent system (carotenoids), 495
 — solvent system (carotenoids, total lipids), 530
 — solvent system (fatty acids, phosphatides), 519
 — solvent system (glycerides; phosphatides), 228
 — solvent system (inosine), 184
 — solvent system (lipid mixtures), 51
 — solvent system (myelin lipids), 157
 — solvent system (oxy-fatty acids), 489
 — solvent system (phospholipids), 101, 107, 111, 187
 — solvent system (phytanic acid glycerol cholesterol esters), 389

- solvent system (sterols), 239
 - solvent system (triglycerides), 30
 - sterols: cholesterol, stigmasterol, β -sitosterol (silica gel HF), 239
 - triglycerides, 170
 - triglycerides (silica gel G-silver nitrate), 30
 - triglycerides by number of double bonds (silica gel-silver nitrate), 539
 - triglycerides by number of double bonds (silica gel G-silver nitrate), 291
 - two-dimensional for liver lipids, 287
 - two-dimensional (phospholipids), 187, 267
 - two-dimensional, phospholipids (silica gel H-magnesium silicate, 101
 - Chromium trioxide, hydroxy fatty acids oxidation, 65
 - Chylomicrons, lack of electrophoretic migration in a starch gel, 420
 - Chymotrypsin-A EC-3.4.4.5, effect of lipids, 221
 - Claviceps species, fatty acid profiles, 260
 - oxygen containing fatty acids, 260
 - Claviceps sulcata, (+)-threo-9,10-dihydroxystearic acid, 260
 - Clostridiopeptidase EC-3.4.4.19, effect of lipids, 221
 - Clover. See plants
 - CoA. See coenzyme A
 - Cockroach, American. See insects
 - Cod, isomer distribution in monoenoic esters, 313
 - triglycerides, positional distribution of fatty acids, 24
 - triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - Cod liver, isomer distribution in monoenoic esters, 313
 - Cod liver oil, autooxidation, 88
 - Cofactors, for α -oxidation, 275
 - Colostrum, human, triglyceride, cholesterol and fatty acid profile, 471
 - Columba livia. See pigeon
 - Column chromatography. See chromatography, column
 - Column selection for GLC, evaluation, 374
 - Combustion of tissues, apparatus and method, 225
 - for radio isotope dtm 225
 - Conidiobolus denesporus. See microorganisms
 - Contaminants, 3,5-di-tert-4-hydroxytoluene in anhydrous diethyl ether, 557
 - sources, Negal plastic, 239
 - sources, polypropylene, 192
 - Contamination, of water by feces 301
 - 5 β -Coprostan-3 β -ol, GLC conditions and derivatives for GLC, 458
 - Coprosterol. See 5 β -cholestan-3 β -ol
 - Cordia salicifolia Cham., fatty acid profile, properties, 43
 - Cordia verbenacea DC., fatty acid profile, properties, 43
 - Cordiidae, fatty acid profile, properties, 43
 - Coriolic acid, (R)-13-hydroxy-cis-9,trans-11-octadecadienoic acid, 425
 - isolation, structure, positional distribution in triglycerides, possible biogenetic significance, 425
 - from sunflower seeds, 489
 - See also (R)-13-hydroxy-cis-9,trans-11-octadecadienoic acid
 - Cormorant, triglycerides, positional distribution of fatty acids, 24
 - Coronarinic acid, from sunflower seeds, 489
 - See cis-9-L,10-L-epoxy-cis-12-octadecenoic acid
 - Cotton plant tissue, 495
 - Cottonseed, isomer distribution in monoenoic esters 313
 - Countercurrent distribution (CCD), carotenoids of potato, 530
 - fatty acid methyl esters, 37
 - Crepennic acid. See cis-9-octadecen-12-ynoic acid
 - Crepis rubra, seed oil comp and production, 307
 - Cricket, house. See insect, Acheta domesticus L.
 - Crotalus adamanteus, phospholipase activity, 107
 - Crotalus atrox. See phospholipase A
 - Cryptantha barbiger (Gray) Greene, fatty acid profile, properties, 43
 - Cyanogen iodide, lipid antioxidant property, 182
 - Cynoglossace, fatty acid profile, properties, 43
 - Cynoglossum creticum Mill., fatty acid profile, properties, 43
 - Cynoglossum lanceolatum Forsk., fatty acid profile, properties, 43
- D**
- Decarboxylation, of α -hydroxy acids, 275
 - of keto acids, 275
 - 4-Decenoic acid (10:1 ω 6), methyl ester, synthesis, properties, metabolism in rats, 14
 - Decyl sulfuric acid, sodium salt, use in solubilization of succinylated apoprotein from β -lipoprotein, 463
 - 4-Decynoic acid, methyl ester, synthesis, properties, 14
 - synthesis, properties, 14
 - LIPIDS, VOL. 3, NO. 6
 - 7-Dehydrocholesterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - 24-Dehydrocholesterol. See desmosterol
 - 7-Dehydrocholesterol trimethylsilyl ethers, mass spectrum, 391
 - Dehydroepiandrosterone, GLC, 374
 - Dehydrogenases, inhibition by oxidized phosphatidic acid, 456
 - Desaturation of fatty acids, effect of temp, 121
 - Desmosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - mass spectrum, 391
 - See also 24-dehydrocholesterol
 - Desmosterol acetate, mass spectrum, 391
 - Desmosteryl trimethylsilyl ethers, mass spectrum, 391
 - Detergent, effect on lipid adsorption, 439
 - Development, insects, neutral lipids and fatty acid profile, 247
 - Dextran gel columns, elution of proteolipids, 453
 - Diadinoxanthin, of *Gyrodinium resplendens*, 5
 - possible identity with antheraxanthin, 5
 - spectra, ultra-violet, visible, 5
 - spectra, ultra-violet, visible; column chromatography, 5
 - Diadinoxanthin-furanoid-oxide, spectra, ultra-violet, visible; column chromatography, 5
 - Dietary fat, fatty acid profile, comparison with human milk, 471
 - Diets, calcium effect on rat hypolipemia, 1
 - effects on bile acids comp. of sheep and bovine, 72
 - fat, rat, effect on metabolism of acetoacetic acid, 545
 - human, comp, 147
 - rat, cocoa butter, effect of strain variety, 1
 - rat, corn oil, effect of strain variety, 1
 - rat, corn-soya, effect of strain variety 1
 - rat, effect on intestinal silicic metabolism, 346
 - for sheep and bovine, synthetic and natural, 72
 - Diffusion, immuno-, human serum β -lipoprotein, 463
 - Digalactosyl diglyceride, cotton buds, fatty acid profile, 373
 - fatty acid profile, in *Spirulina platensis* (algae), 46
 - in maturing wheat, 557
 - from potato tubers, 477
 - Diglyceride kinase, use in dtm of positional distribution of fatty acids in triglycerides, 30
 - Diglycerides, hydrolysis by *Vernonia anthelmintica* lipase, 498
 - incorporation of 14 C-glycerol, 411
 - Dihydrogenosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - Dihydrocholesterol. See 5 β -cholestan-3 β -ol
 - Dihydroxy acid, acetate esters, *Cardamine impatiens* triglycerides, 215
 - 3-keto-7 α ,12 α -Dihydroxy-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,7 α -Dihydroxy-12-keto-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,7 α -Dihydroxy-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,7 β -Dihydroxy-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,12 α -Dihydroxy-7-keto-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,12 α -Dihydroxy-5 β -choleanoic acid, in bile, sheep, fetus; GLC, 72
 - 3 α ,12 β -Dihydroxy-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,7 α ,12 α -Dihydroxy-5 β -choleanoic acid in bile, sheep, bovine, fetus; GLC, 72
 - 3 α ,7,12-Dihydroxy-5 β -choleanoic acid in bile, sheep, bovine, fetus; GLC, 72
 - 3 β ,12 α -Dihydroxy-5 β -choleanoic acid, in bile, sheep, bovine, fetus; GLC, 72
 - 3,10-Dihydroxydecanoic acid, IR and optical rotatory dispersion, spectra, 535
 - from royal jelly, 535
 - Dihydroxystearic acid, in *Claviceps* species, 260
 - (+)-threo-9,10-Dihydroxystearic acid, in *Claviceps sulcata*, 260
 - Diimide reduction, of unsaturated fatty acid, 489
 - Dilinoleoylglycerophosphoryl ethanolamine. See phosphatidyl ethanolamine
 - 4,4-Dimethylcholesterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - N,N-Dimethyl phosphatidyl ethanolamine, IR identification, 384
 - IR spectrum, 104
 - Dimethylpolysiloxane polymer, for GLC, 374
 - phenyl substituted, for GLC, 374
 - 6,7-Dimethyl-5,6,7,8-tetrahydropterine, cofactor for α -oxidation, 275
 - 5,7-Dimethyl tocol, *in vivo* antioxidant, 331
 - 7,8-Dimethyl tocol (γ -tocol), *in vivo* antioxidant 331
 - 4,4-Dimethyl trimethylsilyl ethers, mass spectrum, 391

- Dinoxanthin, of *Gyrodinium resplendens*, 5
— spectra, ultra-violet, visible, 5
— spectra, ultra-violet, visible; column chromatography, 5
Dinoxanthin-furanoid-oxide, spectra, ultra-violet, visible; column chromatography, 5
rac-1,2-Diolefin, synthesis, 228
1,2-Dioleoyl glycerol-3-phosphoryl-hydroxy-L-proline. See phosphatidyl hydroxy-L-proline
Dioleoylglycerophosphoryl-N-anisoyloxycarbonyl-hydroxy-L-proline, phthalimidomethyl ester, synthesis and properties, 228
Dioleoylglycerophosphoryl ethanolamine. See phosphatidyl ethanolamine
Dioleoylglyceryl(2-aminoethyl)phosphonate, activity in blood clotting, 234
Dioleoylglyceryl(2-aminoethyl)phosphonate, total synthesis, 234
Dioleoyl glyceryl(2-phthalimidomethyl)-phosphonate, synthesis and properties, 234
rac-Dioleoyl 3-glyceryl(2-aminoethyl)-phosphonate, synthesis and properties, 234
Dioleoyl phosphatidyl hydroxy-L-proline. See phosphatidyl hydroxy-L-proline
Diphenylamine inhibition, of β -carotene biosynthesis in *Phycomyces*, 558
Diphosphatidyl glycerol. See cardiolipin
Docosahexaenoic acid (22:6), positional distribution in triglycerides, 417
Docosahexaenoic acid (22:6 ω 3), autooxidation, 88
Docosapentaenoic acid (22:5), positional distribution in triglycerides, 417
Docosapentaenoic acid (22:5 ω 3), autooxidation, 88
cis-13-Docosonic acid (22:1), methyl ester, NMR spectrum, 193
Dodeca-3,6-dienoic acid (12:2 ω 6), methyl ester, synthesis, properties, metabolism in rats, 14
Dodeca-3,6-diene-1-ol, synthesis, properties, 14
Dodeca-3,6-dienoic acid, synthesis, properties, 14
Dog, hyperlipemia, effect on circulating leukocytes and erythrocytes, 84
— hyperlipemia, experimental endogenous, 84
— isomer distribution in monoenoic esters, 313
Dog adipose tissue, triglyceride profile, 539
Dog inguinal subcutaneous tissue, triglyceride profiles, 539
Dog mesenteric adipose tissue, triglyceride profiles, 539
Dog pericardial adipose tissue, triglyceride profiles, 539
Dog perirenal adipose tissue, triglyceride profiles, 539
Domestic sewage, sterol content, 301
Double bond determination, by ozonolysis, 403
Double bond position, dtm by periodic-permanganate oxidation, 96
Douglas fir. See trees, also seeds
Doxantha unguis-cati, isomer distribution in monoenoic esters, 313
Duck, isomer distribution in monoenoic esters, 313

E

- EC 1.1.1.8, D-glucose-6-phosphate = NAD oxidoreductase, 456
— L-glycerol-3-phosphate:NAD oxidoreductase, 456
EC 2.7.1.40. See pyruvic acid kinase
EC 3.1.1.4. See phospholipase A
EC 3.1.2.2. See palmitoyl-coenzyme A hydrolase
EC 3.1.3.2. See phosphatase, acid
EC 3.1.3.4. See L- α -phosphatidate phosphohydrolase
EC 3.1.6.1. See arylsulfatase
EC 3.2.1.36. See β -acetylglucosaminidase
EC 3.4.4.1. See pepsin
EC 3.4.4.4. See trypsin
EC 3.4.4.5. See chymotrypsin-A
EC 3.4.4.10. See papain
EC 3.4.4.12. See ficin
EC 3.4.4.19. See clostripain
EC 3.4.4.23. See cathepsin-D
EC 3.4.4.24. See bromelain
Echiaeae, fatty acid profile, properties, 43
Echium rubrum Jacq., fatty acid profile, properties, 43
Echium vulgare L., fatty acid profile, properties, 43
EFA. See fatty acids, unsaturated
cis-5-Eicosanoic acid (20:1 ω 5), in *Caltha palustris* L., 37
cis-11-Eicosanoic acid (20:1 ω 9), in *Caltha palustris* L., 37
Eicosapentaenoic acid (20:5 ω 3), autooxidation, 88
Eicosatetraenoic acid, from *Conidiobolus denacapsus*, 368
— in *Ephedra nevadensis* seed fat, 170
cis,cis,cis,cis-5,11,14,17-Eicosatetraenoic acid (20:4 ω 3), in *Caltha palustris* L., 87
cis,cis,cis,cis-5,11,14,17-Eicosatetraenoic acid (20:4 ω 3) methyl ester, NMR spectrum, 37

- Eicosatrienoic acid, in *Ephedra nevadensis* seed fat, 170
— influence of sex hormones on EFA deficiency in rats, 199
cis,cis,cis-5,11,14-Eicosatrienoic acid (20:3 ω 6), in *Caltha palustris* L., 37
Elaidic acid (18:1), methyl ester, NMR spectrum, 193
Electron micrograph, β -apoprotein from β -lipoprotein, 463
Electron microscopy, of subcellular fractions, 482
Electrophoresis, immuno-, human serum β -lipoprotein, 463
— starch gel, human serum β -1-lipoprotein, 463
— starch gel, serum α -1-lipoproteins, reproducibility, staining, 420
Embryological development, rabbit, fatty acid profiles, lipids synthesis, 361
Endometrium, human, phospholipid comp, 349
— phospholipid incorporation of acetate in vivo, 349
— rabbit, phospholipid comp, 349
Entomophthoraceae. See microorganisms
Enzyme, inhibition by lipid, 456
— inhibition by retinol, 221
— involvement in acetylenic fatty acid synthesis in *Crepis rubra*, 307
— kinetic data, phospholipid activator, 111
Ephedra nevadensis seed fat, fatty acid profile, 170
— triglyceride profile, 170
Epinephrine, effect on 4-¹⁴C-cholesterol distribution in dogs, 324
Epoxide detection, 5
Epoxy. See name of parent compound
Epoxy acids, synthesis, oxygen requirement, 163
9,10-Epoxyoctadecanoic acid, biosynthesis by *Puccinia graminis* (Pers.), 163
cis-9,10-Epoxy-cis-12-octadecenoic acid, from sunflower seeds, 489
cis-9,10-Epoxy-cis-12-octadecenoic acid (18:1 ω 6), *Aster alpinus*, oilseed, 81
cis-9,10-Epoxyoctadecanoic acid, from sunflower seeds, 489
Epistemicus fuscus. See bat
Equations, for computation of brain lipid comp, 284
Ergosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Ergosterol trimethylsilyl ethers, mass spectrum, 391
Eritrichiaeae, fatty acid profile, properties, 43
Erucic acid (22:1), methyl ester, NMR spectrum, 193
Erythrocytes, dog, cholesterol, phospholipids, triglycerides in Triton hyperlipemia, 84
— fatty acid profiles in Triton hyperlipemia, dog, 84
— high potassium cells and low potassium cells, phospholipid comp, 267
— human, fatty acids, 387
— sheep, phospholipid comp, 267
Essential fatty acid deficiency, in rat, 199
Essential fatty acids, deficiency in rats, effect on liver mitochondria swelling and phospholipase A₂, 449
Esters. See also specific fatty acids
Estradiol, influence on lipids, influence of sex hormones on EFA deficiency in rats, 199
Estrogens, GLC column selection, 374
Ethanolamine plasmalogen, rat brain myelin development, 157
Ether phospholipids, biosynthesis from glyceryl ethers, 51
Ethers, alk-1-enyl glyceryl, hydrocarbon profile, of insects, 503
— alkyl glyceryl, hydrocarbon profile, of insects, 503
— alkyl glyceryl, rat metabolism of, 51
— diethyl, BHT as an artifact from, 557
— glycerol, of insects, 503
— glyceryl, solution with sodium taurocholate, 51
— glycerol lipid, enzyme hydrolysis, 111
— 1,2-propanediol 3-hexadecyl. See α -chimy alcohol
— 1,2-propanediol 3-octadecyl. See α -batyl alcohol
— 1,3-propanediol 2-hexadecyl. See β -chimy alcohol
— 1,3-propanediol 2-octadecyl. See β -batyl alcohol
— See also parent compound
Etiocanolone, GLC, 374
Euglena gracilis. See microorganisms

F

- Fasted animals. See starvation
Fat body, glyceryl ether content in insects, 503
Fat deficient diet, rat, supplemented by experimental lipid comps or linoleic acid, 14
Fat diet, rat, effect on metabolism of acetoacetic acid, 545
Fats, animal, cis-9-hexadecenoic and cis-11-octadecenoic acid content of, 313
— bovine milk (triglyceride profile), 291
— dietary, fatty acid profile, comparison with human milk, 471
Fatty acid oxidation, by bat brown adipose tissue, temp effect, 840

- Fatty acids, acetylenic, biosynthesis by *Crepis rubra*, 307**
 — in *Acheta domesticus*, 247
 — autooxidation of saturated, 59
 — biosynthesis of, in rat skin surface lipid, 403
 — branched chain, in *Conidiobolus denaesporus*, 368
 — butyl esters, GLC, 291
 — chain elongations of, 403
 — comp in ass milk phospholipids, 107
 — comp in camel milk phospholipids, 107
 — comp in pig milk phospholipids, 107
 — comp, rust-infected wheat, 163
 — conjugated trienoic, biogenesis, 425
 — desaturation mechanisms of, 403
 — (+)-*threo*-9,10-dihydroxystearic acid in *Claviceps sulcata*, 260
 — epoxy, from sunflower seed oils, 489
 — essential fatty acid deficiency in rats, 449
 — of fish, effect of temp, 121
 — fungal glycerides, acetate and glucose incorporation, 211
 — fungal spore phospholipids, 459
 — fungal spores, relation to accumulation of chemicals, 459
 — GLC of silyl esters, 425
 — of human erythrocytes phospholipids, 387
 — human pancreas lipids comp, 321
 — hydroxydienic, positional distribution, possible biogenetic significance, 425
 — keto, decarboxylation, 275
 — losses during saponification extraction of small samples, 189
 — metabolism of adipose tissue, rat, 383
 — metabolism, chloroplast lipids as intermediates in, 354
 — metabolism in green and blue-green algae, 354
 — metabolism in rust-infected wheat, 163
 — methyl ester, diacyl, GLC, 65
 — methyl esters, GLC column selection, 374
 — methyl esters, in human pancreas lipids, 321
 — methyl ester, monobasic, GLC, 65
 — methyl esters GLC, column chromatography, 403
 — methyl esters in grasshopper eggs (*Melanoplus bivitatus*), 455
 — NMR spectrum of double bond protons in aliphatic systems, 193
 — non-esterified, human brain, equations for computation, 284
 — non-esterified, in human pancreas, 321
 — odd and branched chains, 403
 — oxidation by brown adipose tissue homogenates, 340
 — oxidation of saturated, 59
 — pollen lipids attractive to honey bees, 530
 — positional distribution of fatty acids in rat liver, 30
 — positional distribution in phosphatidyl choline, 373
 — positional distribution in triglycerides of aquatic animals, 24, 417
 — positional distribution in triglycerides, formula for estimation pattern, 24
 — positional distribution in triglycerides, prediction by theory, 417
 — in rat diet, effect on acetate use, 346
 — saponification, losses of, 189
 — saponification, separation of products, 189
 — solution with sodium taurocholate, 51
 — stable pool in adipose tissue, human, 147
 — stereospecific hydrolysis from triglycerides, 519
 — synthesis, relationship between *cis*-9-hexadecenoic and *cis*-11-octadecenoic acid in, 313
 — synthesis, rust-infected wheat, 163
 — tetradecan-1,14-dioic acid, 186
 — utilization for synthesis of diglycerides and triglycerides, 411
 — very long chain, 403
 — Xanthophyll esters in *Tagetes erecta*, 183
- Fatty acids, epoxy, biosynthesis, by *Puccinia graminis* (Pers.), 163**
 — in *Claviceps* species, 260
- Fatty acids, hydroxy, acetoxyhydroxy acid, methyl ester, IR, NMR, mass, 215**
 — in *Claviceps* species, 260
 — decarboxylation, 275
 — mass spectra, 431
 — monoacetate ester, dihydroxydocosanoic acid, 215
 — monoacetate ester, dihydroxyeicosanoic acid, 215
 — monoacetate ester, dihydroxystearic acid, 215
 — monoacetate ester, dihydroxytetraecanoic acid, 215
- Fatty acids, profile, of *Anabaena cylindrica* (algae), 46**
 — of *Anabaena flos-aquae* (algae), 46
 — of *Anacystis nidulans* (algae), 46
 — of ass milk phospholipids, 107
 — of *Aster alpinus* and *Arctium minus* oils, 491
 — of autooxidized cod liver oil, 88
 — bovine milk triglycerides, 291
 — of brain glucosyl ceramide, 262
 — *Caltha palustris* L., 37
 — of camel milk phospholipids, 107
 — of *Chlorogloea fritschii* (algae), 46
 — of *Claviceps* species, 260
 — of *Conidiobolus denaesporus*, 368
 — di- and triglycerides, 411
 — of dog erythrocytes in Triton hyperlipemia, 84
 — of fish, effect of temp, 121
 — fungal spore phospholipids, 459
 — of glyceryl ethers, 511
 — of goldfish phosphatidyl choline, 121
 — of goldfish phospholipid, 121
 — of goldfish sphingomyelin, 121
 — of goldfish total lipid, 121
 — of goldfish triglyceride, 121
 — of house cricket (*Acheta domesticus* L.), 247
 — human adipose tissue, 147
 — human endometrium phospholipids, 349
 — human pancreas cholesterol esters, 321
 — human pancreas lipids, 321
 — human pancreas nonesterified fatty acids, 321
 — human pancreas phospholipids, 321
 — human pancreas triglycerides, 321
 — human skin surface lipid, 403
 — of lipids of potato tubers, 477
 — of liver, rats fed experimental fatty acids, 14
 — of *Mastigocladus laminosus* (algae), 46
 — of maturing wheat, 557
 — of methyl esters in grasshopper eggs, 455
 — of mouse brain phospholipids, 79
 — of *Myxosarcina chroococcoides* (algae), 46
 — of neutral lipids and glycerides of *Spirulina platensis* (algae), 46
 — in phospholipids from camel, ass, and pig milk, 107
 — of pig milk phospholipids, 107
 — polar lipids of bud of cotton plants, *Gossypium* sp., 373
 — of *Pseudomonas aeruginosa* grown in hexadecane, 460
 — of *Puccinia graminis* (Pers.), 163
 — monore with unusual double bond positions, 403
 — monoenic isomers, distribution in rat skin, 403
 — rabbit endometrium phospholipids, 349
 — rabbit liver, embryological development, 361
 — rat liver, 30
 — rat serum total lipids, effect of diet, 545
 — rat skin surface lipid, 403
 — from seed oils. See oilseeds
 — of *Spirulina platensis* (algae), 46
 — from sunflower seed oils, 489
 — of triglycerides in human colostrum and milk, 471
 — of wheat rust, 163
- Fatty acids, unsaturated, biosynthesis by *Crepis rubra*, 307**
 — circum-annual changes in brown adipose comp, 340
 — diimide reduction of, 489
 — dtmn of double bond position, 379
 — dtmn double bond position of monoenoic acid, 96
 — of fish, 121
 — liver, rats fed experimental fatty acids, 14
 — NMR spectra, 193
 — nonketogenic in rats, 545
 — pollen lipids attractive to honey bees, 519
 — positional distribution in triglycerides of aquatic animals, 24
 — rats fed experimental fatty acids, 14
 — *cis*-5-unsaturated in *Caltha palustris* L., 37
- Fatty acid, unsaturated essential, deficiency in rats, 199**
- Fatty acids, unsaturated, poly-, arachidonic, EFA deficiency, 199**
 — deficiency in rats, 199
 — docosahexaenoic acid, positional distribution in triglycerides, 417
 — docosapentaenoic acid, positional distribution in triglycerides, 417
 — eicosatrienoic, EFA deficiency, 199
 — formation in fish, 121
 — mobilization of linoleic, 199
 — positional distribution in triglycerides of aquatic animals, 24
 — in rat diet, effect on acetate use, 346
 — *cis*-5-unsaturated in *Caltha palustris* L., 37
- Fatty aldehydes, profile, of mouse brain phospholipids, 79**
- Fecal lipids, rat, in rat hypolipemia, effect of strain variety, 1**
- Feces, bovine, bile acids, comp, 72**
- Ferrous ion, cofactor for α -oxidation, 275**
- Fetus bile acids, sheep, bovine, 72**
- Ficin EC-3.4.4.12, effect of lipids, 221**
- Fir, Douglas. See trees, also seeds**
- Fish, fresh water, triglycerides, positional distribution of fatty acids, 24**
 — *Heteropneustes fossilis*, phospholipids of tissues, 21
 — marine, triglycerides, positional distribution of fatty acids, 24

- triglycerides, positional distribution of docosapenta-**Goldfish**, temp on fatty acid profile, 121
 - See specific name
 - Fish and invertebrates**, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - Flavoxanthin**, from cotton plants, properties, 495
 - Frog**, triglycerides, positional distribution of fatty acids, 24
 - Frog brain**, ganglioside biosynthesis, *in vitro*, 378
 - Fruit**, See oils, seeds, oilseeds or specific name
 - Fucosanthin**, in a marine dinoflagellate, 5
 - Fungal glycerides**, acetate and glucose incorporation, 211
 - Fungus**, See microorganisms
- G**
- Galactolipid**, cholesterol complex, 157
 - in potato tubers, 477
 - rat brain myelin development, 157
 - See galactosyl diglyceride
 - Galactose**, incorporation into glycosphingolipids, 151
 - Galactosyl ceramide**, bovine brain, IR, 262
 - Galactosyl diglyceride**, cotton buds, fatty acid profile, 373
 - fatty acid profile in *Spirulina platensis* (algae), 46
 - in maturing wheat, 557
 - metabolism in green and blue-green algae, 354
 - Ganglioside biosynthesis**, frog brain, 378
 - Gangliosides**, brain glucosyl ceramide, as a precursor of, 262
 - human brain, equations for computation, 284
 - Ganglioside (monosialo)**, formation enzymatically from (Tay-Sachs) ganglioside, 378
 - Ganglioside (Tay-Sachs)**, conversion to (monosialo) ganglioside by adult frog brain homogenate, 378
 - Gas-liquid chromatography**, See chromatography, gas-liquid
 - Gastrocyle hispidus Bunge**, fatty acid profile, properties, 43
 - Geotrichum candidum**, See microorganisms, also lipase
 - Germination**, of seeds, Douglas fir, 482
 - Glomerella cingulata**, glycerides, relative incorporation of precursors, 211
 - Glucose**, incorporation into fungal glycerides, 211
 - incorporation into glycosphingolipids, 151
 - Glucose-¹⁴C and galactose-¹⁴C conversion**, to glycosphingolipids, 151
 - Glucose incorporation**, into brain glycolipids, 262
 - Glucose-6-phosphate dehydrogenase EC 1.1.1.49**, lipid inhibitor, 456
 - D-Glucose-6-phosphate: NADP oxidoreductase EC 1.1.1.49**, lipid inhibitor, 456
 - Glucosyl ceramide**, bovine brain, IR, 262
 - in human leukocytes, 151
 - spleen from Gaucher's disease, 262
 - Glycerides**, in bovine milk, 291
 - in crickets, 247
 - fungus, acetate and glucose incorporation, 211
 - influence of sex hormones on EFA deficiency in rats, 199
 - in maturing wheat, 557
 - phytanic acid, TLC, 389
 - phytanic acid containing, prep, 389
 - Glycerol**, fungal glycerides, acetate and glucose incorporation, 211
 - release from rat adipose tissue, PGE, effect, 383
 - ¹⁴C-Glycerol**, incorporation into diglycerides and triglycerides, 411
 - Glycerol ether ozonides**, reduction with dimethyl sulfide, 511
 - reduction with lithium aluminum hydride, 511
 - n-Glycerolphosphate dehydrogenase EC 1.1.1.8**, lipid inhibitor, 456
 - L-Glycerol-3-phosphate=NAD oxidoreductase EC 1.1.1.8**, lipid inhibitor, 456
 - Glycerolphospholipid**, in human organs, accumulation, 287
 - Glycerol alkyl ethers**, rat metabolism of, 51
 - Glycerol ethers**, acetoxy-mercuri-methoxy derivatives, separation of, 511
 - alkyl iodides, conversion to, anal of, 511
 - biosynthesis, from ¹⁴C-1-acetate in insects, 503
 - changes during insect development, 503
 - content in insects, 503
 - conversion to allyl alkyl ethers and their preparative GLC, 511
 - reductive ozonolysis, location of double bonds, 511
 - solution with sodium taurocholate, 51
 - TLC, 129
 - Glycerol hexadecyl ether**, See chimyl alcohol
 - Glycerol octadecyl ether**, See batyl alcohol
 - Glycocholic acid**, sodium salt, effect on aortic hydrolysis of cholesteryl oleate, 454
 - Glycolipids**, of bovine brain, 262
 - Glycosphingolipid**, biosynthesis in human leukocytes, 151
 - See glycolipid, sphingolipid, sphingosine
- H**
- Hackelia jessicae (McGreg.) Brand**, fatty acid profile, properties, 43
 - Harbor seal**, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
 - triglycerides, positional distribution of fatty acids, 24
 - Harpagonellase**, fatty acid profile, properties, 43
 - Harp seal**, triglycerides, positional distribution of fatty acids, 24
 - triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
 - HDL**, See lipoprotein, high density
 - See high density lipoproteins
 - Heart**, human coronary disease, adipose tissue, 147
 - fish, (*Heteropneustes fossilis*), phospholipids, 21
 - pigeon, (*Columba livia*), phospholipids, 21
 - rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
 - toad, (*Bufo melanostictus*), phospholipids, 21
 - turtle, (*Kachuga Smithi*), phospholipids, 21
 - Heart atrial myocardium**, dog, distribution between tissue and plasma, 324
 - Heart ventricular myocardium**, dog, distribution between tissue and plasma, 324
 - Heliothis virescens**, See insects
 - Heliotropioideae**, fatty acid profile, properties, 43
 - Heliotropium amplexicaule Vahl**, fatty acid profile, properties, 43
 - Heliotropium curassavicum L.**, fatty acid profile, properties, 43
 - Heliotropium strigosum Willd.**, fatty acid profile, properties, 43
 - Heliotropium supinum L.**, fatty acid profile, properties, 43
 - Hepatic**, See liver
 - n-Hept-2-enal**, product of autooxidation, 88
 - Hereditary metabolic disorders**, lipidosis, 287
 - Herring**, triglycerides, positional distribution of fatty acids, 24
 - Herring gull**, triglycerides, positional distribution of fatty acids, 24
 - Herring (Icelandic)**, isomer distribution in monoenoic esters, 313
 - Heterokaryons**, of *Physcomyces* mutants, biosynthesis of β -carotene, 558
 - Heteropneustes fossilis**, See fish
 - Hexadecenyl glycerols**, double bond position dtmn, 511
 - trans-3-Hexadecenoic acid (16:1c3)** *Aster alpinus*, oil-seed, 91
 - cis-3-Hexadecenoic acid**, in natural fats and oils, 313
 - 1-Hexadecyl glyceryl ether**, See α -chimyl alcohol
 - 2-Hexadecyl glyceryl ether**, See β -chimyl alcohol
 - Hexadeca-7,10-dienoic acid (16:2c6)**, methyl ester, synthesis, properties, metabolism in rats, 14
 - Hexadeca-7,10-divenoic acid**, synthesis, properties, 14
 - n-Hex-2-enal**, product of autooxidation, 88
 - 5-Hexynoic acid**, synthesis, properties, 14
 - Hibernation**, bat, circum-annual variation of brown adipose tissue triglycerides, 340
 - High density lipoproteins**, human, electrophoresis in a starch gel, 420
 - Hippopotamus**, isomer distribution in monoenoic esters, 313
 - Honey bees**, pollen lipids attractive to, 530
 - Honey bees (*Apis mellifera* L.)**, See insects
 - Human**, adipose tissue, fatty acid turnover, pool size, 147
 - coronary heart disease adipose tissue, 147
 - diet comp, 147
 - fatty acid profile, adipose tissue, 147
 - fatty acids of skin surface lipid, 403
 - leucocytes, sphingolipid comp, 151
 - Human brain**, annual change in comp, 284
 - lipid comp, age changes, 284
 - Human colostrum**, triglyceride, cholesterol and fatty acids profile, 471
 - Human endometrium**, phospholipid comp, acetic acid incorporation, 349

- Human liver**, normal comp, 287
 — sphingomyelinosis, 287
Human milk, phospholipids profile, 101
 — triglyceride, cholesterol and fatty acids profile, 471
Human pancreas, lipid comp, 321
Human plasma lipids, Refsum's syndrome, diagnosis by TLC, 389
Human serum, lipoprotein, succinylated, 463
Human serum, lipoprotein succinylated, 463
 — starch gel electrophoresis, 463
 — variants, α -1-lipoproteins, 420
Hyamine, alkyl (C^2 to C^{25}) tolyl methyl trimethyl ammonium chloride, 225
 — solubilization of tissue for liquid scintillation counting, 225
Hydrazine reduction, of unsaturated fatty acids, 37
Hydrocarbons, insects, *Acheta domesticus*, 250
Hydrogen-3. See tritium
Hydrogenation, micro, and GLC of hydroxydienoid fatty esters, 425
Hydrogenolysis, lithium aluminum hydride reduction, 129
Hydrolase, alkenyl glycerol-3-phosphoryl choline, 111
Hydroperoxides, from saturated fatty acids autooxidation, 65
 α -Hydroxy acids, decarboxylation by brain microsomes, 275
3-keto-12-Hydroxy-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
3 α -Hydroxy-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
3 α , Hydroxy-7-keto-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
3 α , Hydroxy-12-keto-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
3 β -Hydroxy-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
7-Hydroxycholesterols, synthesis, isolation, properties, GLC, IR, 551
3-Hydroxydecanoic acid, from royal jelly, 535
11- β -Hydroxyetiocholanolone, GLC, 374
3 β -Hydroxy-12-keto-5 β -cholanolic acid, in bile, sheep, bovine, fetus; GLC, 72
9-Hydroxy-trans-10,cis-12-octadecadienoic acid, from sunflower seeds, 489
13-Hydroxy-cis-9,trans-11-octadecadienoic acid, from sunflower seeds, 489
(R)-13-Hydroxy-cis-9,trans-11-octadecadienoic acid, isolation, positional distribution in triglycerides, possible biogenetic significance, 425
 — see coriolic acid
12-Hydroxy-cis-9-octadecenoic acid. See also rincinoic acid
8-Hydroxyoctanoic acid, from royal jelly, 535
Hydroxy palmitic acid, methyl ester, oxidation by chromium trioxide, 65
Hypercholesterolemia, effect on serum phosphatidyl choline-cholesterol acyltransferase, 381
Hyperlipemia, dog, effect on circulating leukocytes and erythrocytes, 84
 — experimental endogenous, dogs, 84
 — rat, effect of dietary calcium, 1
 — rat, Holtzman variety, 1
 — rat, strain differences, diet, 1
 — rat, Wistar variety, 1
 — Triton, effect on serum phosphatidyl choline-cholesterol acyltransferase, 381
 — Triton WR-1339 induced, 84
- I**
- Imidazole**, inhibition of alkenyl hydrolase, microsomes, 111
Immunodiffusion. See diffusion, immuno-
Immuno-electrophoresis. See electrophoresis, immuno-
Indian buffalo milk, phospholipids profile, 101
Inguinal subcutaneous adipose tissue, dog, triglyceride profiles, 539
Inosine, in lipid extracts, removal by Sephadex chromatography, 184
 — TLC, 184
Insect development, changes in glyceryl ethers during, 503
 — neutral lipids and fatty acid profile, 247
Insect lipids, anal of glyceryl ethers by TLC and GLC, 503
 — chain length of glyceryl ether side chains, 503
 — fatty acid methyl esters in grasshopper eggs, 455
 — fatty acid profile in the house cricket, 247
 — glyceryl ether content, 503
Insects, *Acheta domesticus* L., house cricket, 250
 — *Anthonomus grandis* (boll weevil), glyceryl ethers during development, 503
 — *Anthonomus grandis* Boheman (boll weevil), feeding stimulant of cotton plants, 495
 — *Apis mellifera* L. (honey bee), fatty acids in royal jelly, 535
 — Comstock mealybug, 186
 — fatty acids and dietary lipids, 247
 — *Heliothis virescens* (tobacco bud worm), glyceryl ethers during development, 503
 — Homoptera: Pseudococcidae, 186
 — hydrocarbons in the common house cricket, 250
 — lipid metabolism, 247
 — *Melanoplus bivitatus*, fatty acid methyl ester profiles of eggs, 455
 — *Periplaneta americana* (American cockroach), glyceryl ethers in whole insect and fat body, during development, 503
 — sterol metabolism, cricket, 256
 — sterols of *Acheta domesticus*, 256
Insect scale, *Pseudococcus comstocki* Kuwana, content of dibasic acid, 186
Intestinal alics, rat, metabolism, effect of diet, 346
Invertebrates, triglycerides, positional distribution of fatty acids, 24
 — See specific name
Iodine, lipid antioxidant property, 182
Iodine compounds, lipid antioxidant property, 182
Iodine cyanide, lipid antioxidant property, 182
Ionel. See 3,5-di-tert-4-hydroxytoluene
Isolulein, from cotton plants, properties, 495
Isomerization, iodine catalyzed, of chloroplast pigments, 5
Isotope, radio active, effect of volume on liquid scintillation efficiency, 225
 — radio active, measurement in tissue of ^3H and ^{14}C , 225
- K**
- Kachuga Smithi**. See turtle
 α -Keto acids, decarboxylation by brain microsomes, 275
7-Ketocholesterol, synthesis, isolation, properties, GLC, IR, 551
Keto compounds. See parent compounds
11-Ketoetiocholanolone, GLC, 374
 α -Ketostearic acids, decarboxylation, 275
17-Keto-steroids, GLC column selection, 374
Kidney, dog, cholesterol distribution between tissue and plasma, 321
 — fish, (*Heteropneustes fossilis*), phospholipids, 21
 — pigeon, (*Columba livia*), phospholipids, 21
 — rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
 — rat, lysosomal cathepsin-D, EC 3.4.4.23, 221
 — toad, (*Bufo malanostictus*), phospholipids, 21
 — turtle, (*Kachuga Smithi*), phospholipids, 21
 K_m values. See enzyme kinetic data
- L**
- Lactosyl ceramide**, in human leukocytes, 151
Lanost-8-en-3 β -ol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Lanosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
Lappula barbatum (Bieb.) Guerke, fatty acid profile, properties, 43
Lard (Hormel), isomer distribution in monoenoic esters, 313
Lauric acid, use in lipoprotein separations, 420
LDL. See lipoprotein, low density
 — See low density lipoproteins
Leucocytes, human, sphingolipid comp, 151
Leukocytes, in Triton hyperlipemia, dog, 84
Lindelofia anchusoides (Lindl.) Lehman, fatty acid profiles, properties, 43
Linoleic acid, in bat brown adipose tissue, 340
 — in crickets, 247
 — effect of polyunsaturated diet, in adipose tissue, human, 147
 — human erythrocytes, other ω -6 fatty acids, 387
 — incorporation into fatty acids of *Crepis rubra*, 307
 — increase in phospholipids during pregnancy, 349
 — methyl ester, oxygen uptake, 182
 — postulated biogenetic conversion to trienoic and hydroxydienoic acids, 425
 — rabbit liver microsomal phospholipids, variation with age, 361
 — in triglyceride, hydrolysis by *Geotrichum candidum*, 143
 — turnover time in adipose tissue stable pool, human, 147
 — use in dietary monitoring with the adipose tissue, human, 147

- in wheat lipids, maturation changes, 557
- Linoleic acid** (18:2 ω 6), carcass fat, influence of sex hormones on EFA deficiency in rats, 199
- methyl ester, NMR spectrum, 193
- mobilization, influence of sex hormones on EFA deficiency in rats, 199
- in potato tubers, 477
- Linolenic acid**, in crickets, 247
- in wheat lipids, maturation changes, 557
- α -**Linolenic acid**. See 9,12,15-octadecatrienoic acid (18:3 ω 3)
- γ -**Linolenic acid**. See 6,9,12-octadecatrienoic acid (18:3 ω 6)
- See *cis,cis,cis*-6,9,12-octadecatrienoic acid
- Linolenic acid** (18:3 ω 3), in potato tubers, 477
- Lipase**, *Geotrichum candidum*, specificity for oleic acid, 143
- *Geotrichum candidum*, stereospecific anal of triglycerides, specificity for *cis*- ω 9, 519
- *Geotrichum candidum*, triglyceride hydrolysis, 143
- of germinating fir seeds, acid, neutral, 482
- pancreatic-EC 3.1.1.3, on dimethyl phosphatides, 191
- pancreatic EC 3.1.1.3, hydrolysis of *Coriaria* oils, 425
- pancreatic, EC 3.1.1.3, use in dtm of positional distribution of fatty acids in triglycerides, 30
- pancreatic, triglyceride stereospecific anal, 519
- *Vernonia anthelmintica* seed, prep of, specificity of, 498
- Lipid composition**, brain, curvilinear regression, 284
- brain, equations for computation, 284
- of wheat, 557
- Lipid extraction**, from algae, 46
- from bile and feces, 72
- from camel, pig and ass milk, 107
- from clover pollen, 530
- from crickets, 247
- from fish (*Heteropneustes fossilis*), 21
- from fungus mycelium, 211
- germinating seeds of Douglas fir, 482
- from *Gyrodinium resplendens*, 5
- from human pancreas, 321
- from human tissues, contamination with inosine, 184
- of mouse brain, 79
- from oilseeds, 37
- from pigeon (*Columba livia*), 21
- from plant tissues, 495
- from rat liver, 14, 30
- from rat liver slices, 411
- from sea water, 239
- from toad (*Bufo melanostictus*), 21
- from tubers, potato, 477
- from turtle (*Kachuga Smithi*), 21
- from wheat, 557
- from wheat rust, 163
- Lipid metabolism**, influence of sex hormones on EFA deficiency in rats, 199
- Lipids**, autooxidation, 88
- of domestic sewage, 301
- fish, effect of temp, 121
- human brain, annual change in comp, 284
- molar ratios of myelin, 157
- phosphatidyl acid-like, from mouse liver, 456
- plasma, Refsum's syndrome, diagnosis by TLC, 389
- in polypropylene, 192
- seed, *Coriaria*, 425
- separation of polystyrene gel columns, 385
- in white potato tubers, 477
- Lipid uptake by membranes**, detergent effect, 439
- effect of salts, 439
- Lipoprotein apoprotein**, from human serum β -lipoprotein, solubilization and properties, 463
- Lipo-: proteins**, aging effects revealed by electrophoresis, 420
- high density, 420
- human β or low density, succinylated apoprotein derivative, 463
- low density, 420
- separation, use of lauric acid, 420
- serum, cholesterol and cholesterol esters in baboon, 136
- serum high density concentration, electrophoretic varieties and conditions for electrophoresis, 420
- serum, incorporation of cholesterol and cholesterol esters, in the baboon, 136
- stain, oil red O, 420
- very low density, 420
- α -1-**Lipoproteins**, staining with oil red O, 420
- β -**Lipoproteins**, human serum, succinylated apoprotein derivative, 463
- staining with oil red O, 420
- Liquid-liquid partition chromatography**. See chromatography, liquid-liquid partition
- Lithospermeae**, fatty acid profile, properties, 43
- Lithospermum arvense** L., fatty acid profile, properties, 43
- Lithospermum purpurascens** L., fatty acid profile, properties, 43

- Liver**, dog, cholesterol distribution tissue and plasma, 324
- fish (*Heteropneustes fossilis*), phospholipids, 21
- mouse, lipid inhibitor of enzymes, 456
- pigeon (*Columba livia*), phospholipids, 21
- rat, lipid content, effect of strain variety, 1
- rat, lipids, experimental diets, 14
- rat, lysosomal cathepsin-D, EC 3.4.4.23, 221
- rat, plasmalogen metabolism, 111
- rat, synthesis, species of diglycerides and triglycerides, 411
- rat, triglycerides, positional distribution of fatty acids, 30
- toad (*Bufo melanostictus*), phospholipids, 21
- turtle (*Kachuga Smithi*), phospholipids, 21
- Liver lipids**, changes in fatty acid profiles with age, rabbit, 361
- rat, influence of sex hormones on EFA deficiency in rats, 199
- Lobster**, isomer distribution in monoenoic esters, 313
- triglycerides, positional distribution of fatty acids, 24
- triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
- Lotus**. See plants
- Low density lipoproteins**, human, migration during starch gel electrophoresis, 420
- human serum, succinylated apoprotein derivative, 463
- Lung**, dog, distribution between tissue and plasma, 324
- Lutein**, from cotton plants, properties, 495
- Lycopodium**, isomer distribution in monoenoic esters, 313
- Lyso bisphosphatidic acid**, sphingomyelinosis, Niemann-Pick disease, 287
- Lysophosphatidyl choline**, in mammalian milk, 101
- TLC, 187
- Lysophosphatidyl ethanolamine**, in mammalian milk, 101
- Lysosomes**, prep rat liver and kidney, 221

M

- Macadamia nut**, isomer distribution in monoenoic esters, 313
- Mackerel**, isomer distribution in monoenoic esters, 313
- triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
- triglycerides, positional distribution of fatty acids, 24
- α -2-**Macroglobulin**, staining with oil red O, 420
- Mammalian milk**, phospholipids profile, 101
- Mammals**. See specific common name
- triglycerides, positional distribution of fatty acids, 24
- Marine dinoflagellate**. See microorganisms, *Gyrodinium resplendens*
- Marine mammals**, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
- triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
- Marine waters**, sterol content, 239
- Marrow**, rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
- Mass spectrometer**. See also chromatography, gas-liquid-mass spectrometer
- Mastigocladus laminosus**, fatty acid profiles, total lipids and neutral lipids, 46
- Mattiasstrum cristatum** (Schreb.) Brand, fatty acid profile, properties, 43
- Maturation**. See development or growth
- Melanoplus bivittatus**, grasshopper. See insects
- Melilotus**. See plants
- Membrane formation**, lipid adsorption in, 439
- Membranes**, adsorption of lipids by, 439
- delipidated, lipid adsorption by, 439
- enzyme-phospholipid interactions, 111
- metabolism of phosphate in phospholipids of erythrocytes, 267
- phospholipids, comp of erythrocytes, 267
- Membranes of fungal spores**, influence on accumulation of chemicals, 459
- Menhaden** (Atlantic), isomer distribution in monoenoic esters, 313
- Menhaden** (Hunko), isomer distribution in monoenoic esters, 313
- Mesenteric adipose tissue**, dog, triglyceride profiles, 539
- Metabolism**, phospholipids, incorporation of phosphate into HK and LK erythrocytes, 267
- Methoxenyl trimethylsilyl ethers**, mass spectrum, 391
- p-Methoxybenzoyloxycarbonyl-hydroxy-L-proline**, synthesis and properties, 228
- Methylation**, of chloroplast pigments, 5
- p-Methylbenzoyloxycarbonyl-hydroxy-L-proline**, phthalimide methyl ester, synthesis and properties, 228
- 4 α -Methyl-5 α -cholest-7-en-3 β -ol**, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391

- 4 β -Methylcholesterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
- 4 β -Methyl cholesterol trimethylsilyl ethers, mass spectrum, 391
- 2-Methyl octacosane, in crickets, 250
- 14-Methyl pentadecanoic acid, methyl ester, from *Conidiobolus denacsporus*, mass spectra, 368
- N-Methyl phosphatidyl ethanolamine, IR identification, 384
- 12-Methyl tetradecanoic acid, methyl ester, from *Conidiobolus denacsporus*, mass spectra, 368
- 8-Methyl tocol (Δ -tocol), in vivo antioxidant, 331
- 12-Methyl tridecanoic acid, methyl ester, from *Conidiobolus denacsporus*, mass spectra, 368
- Mevalonic acid, incorporation into serum cholesterol and cholesterol esters in the baboon, 136
- Microbial disposition, of 5 β -cholestan-3 β -ol, 301
- Microorganisms, *Alternaria olivacea*, 459
- *Claviceps* species, fatty acid comp, 260
 - *Claviceps* species, oxygenated fatty acids, 260
 - *Claviceps* (*ex-Pennicetum*), fatty acid profile, 260
 - *Claviceps gigantea*, fatty acid profile, 260
 - *Claviceps paspali*, fatty acid profile, 260
 - *Claviceps purpurea*, fatty acid profile, 260
 - *Claviceps sulcata*, fatty acid profile, 260
 - *Conidiobolus denacsporus*, fatty acid profile, 368
 - *Euglena gracilis*, 5
 - *Geotrichum candidum*, lipase of, 143
 - *Geotrichum candidum*, lipase for cis- ω 9, 519
 - *Glomerella cingulata*, labeling of glycerides, 211
 - *Glomerella cingulata*, media, 211
 - *Gyrodinium resplendens*, chloroplast pigments, 5
 - *Gyrodinium resplendens*, culture media, 5
 - metabolism of 5 β -cholestan-3 β -ol in sewage, 301
 - *Neurospora sitophila*, 459
- Phycomycetes, inhibition of β -carotene biosynthesis by diphenylamine, 558
- *Phycomycetes* mutants, deficient in β -carotene biosynthesis, 558
 - *Pseudomonas aeruginosa*, fatty acid profile of phosphatidyl ethanolamine, 460
 - *Pseudomonas aeruginosa*, grown in hexadecane, 460
 - *Puccinia graminis* (Pers.), fatty acid comp, synthesis, 163
 - *Rhizopus nigricans*, 459
 - *Spirulina platensis*, fatty acid profiles, trienoic acids, biosynthesis, culture media, 46
- Microsomes, lipase, 482
- phospholipid requirement, 111
 - rat brain, α -oxidation of fatty acids, 275
- Microsomes phosphatidyl choline, changes in fatty acid profiles with age, rabbit, 361
- Milk, ass, phospholipids, fatty acid profiles, 107
- ass, phospholipids profile, 101
 - bovine, phospholipids profile, 101
 - bovine, triglyceride anal, 291
 - bovine, triglyceride structure, 291
 - camel, phospholipids, fatty acid profiles, 107
 - camel, phospholipids profile, 101
 - human, phospholipids profile, 101
 - human, triglyceride, cholesterol and fatty acids profile, 471
 - Indian buffalo, phospholipids profile, 101
 - pig, phospholipids, fatty acid profiles, 107
 - pig, phospholipids profile, 101
 - sheep, phospholipids profile, 101
- Mitochondria, adsorption of emulsified lipids, 439
- lipase, 482
 - rat liver, diet dependence of swelling, 449
 - rat liver, phospholipase A₂ activity, dependency on diet, 449
 - rat liver, swelling change by corn oil diet to essential fatty acid deficient, 449
 - rat liver, swelling in essential fatty acid, 449
- Monoenic fatty acids, dtmn double bond position, 96
- Monogalactosyl diglycerides, from potato tubers, 477
- Monoglycerides, hydrolysis by *Vernonia anthelmintica* lipase, 498
- Mouse brain, phospholipids, fatty acid profile, 79
- phospholipids, fatty aldehyde profile, 79
- Mouse liver, lipid inhibitor of enzymes, 456
- Madpuppy, triglycerides, positional distribution of fatty acids, 24
- Muscle, goldfish fatty acid profile, temp effect, 121
- Muscle, rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
- Mycelium, fungus, acetate and glucose incorporation, 211
- Myelin, adsorption of emulsified lipids, 439
- electron micrograph of isolated, 157
 - molar ratio of lipids, 157
 - protein content during development, 157
 - rat brain development, 157
 - rat brain, prep from young animals, 157

- Myocardium, atrial, dog, distribution between tissue and plasma, 324
- ventricular, dog, distribution between tissue and plasma, 324
- Myristic acid, from *Conidiobolus denacsporus*, 368
- in crickets, 247
- Myxosarcina chroococcoides, fatty acid profiles, total lipids and neutral lipids, 46

N

- NCS, solubilization of tissue for liquid scintillation counting, 225
- Neodiadinanthin, spectra, ultra-violet, visible, 5
- Neodioxanthin, spectra, ultra-violet, visible, 5
- Neoperidin, spectra, ultra-violet, visible, 5
- Neoplastic tissue. See also tumors
- Neoxanthin, from cotton plants, properties, 495
- Neurospora sitophila, fungal spore phospholipids, fatty acid profile, 459
- Neutral lipids, fish, effect of temp, 121
- rat, tissues, 129
- Niemann-Pick disease, accumulation of a glycerolphospholipid, 287
- See sphingomyelinosis
- NMR. See spectra, nuclear magnetic resonance
- n-Nonacosane, in crickets, 250
- Nonnea pulla DC., fatty acid profile, properties, 43

O

- cis-9,cis-12-Octadecadienoic acid (18:2 ω 6), methyl ester, NMR spectrum, 193
- cis,cis,cis,cis-6,9,12,15-Octadecatetraenoic acid (18:4 ω 3), in oilseed, Boraginaceae, 43
- cis,cis,cis-6,9,12-Octadecatrienoic acid (18:3 ω 6), (γ -linolenic acid), in oilseeds, Boraginaceae, 43
- trans-3,cis-9,cis-12-Octadecatrienoic acid (18:3 ω 6), *Arctium minus* and *Aster alpinus* oilseeds, 91
- 6,9,12-Octadecatrienoic acid (18:3 ω 6), in blue-green algae (*Spirulina platensis*), biosynthesis, 46
- by desaturation of linoleic acid, 46
- cis-5-Octadecenoic acid (18:1 ω 13), in *Caltha palustris* L., 37
- cis-9-Octadecenoic acid (18:1), methyl ester, NMR spectrum, 193
- cis-9-Octadecenoic acid (18:1 ω 9), in *Caltha palustris* L., 37
- cis-11-Octadecenoic acid, in natural fats and oils, 313
- trans-3-Octadecenoic acid (18:1 ω 3), *Aster alpinus*, oilseed, 91
- trans-3,cis-9-Octadecenoic acid (18:2 ω 9), *Aster alpinus*, oilseed, 91
- trans-9-Octadecenoic acid (18:1), methyl ester, NMR spectrum, 193
- 1-Octadecyl glyceryl ether. See α -batyl alcohol
- 2-Octadecyl glyceryl ether. See β -batyl alcohol
- Octadecenyl glycerols, double bond position dtmn, 511
- cis-9-Octadecen-12-ynoic acid, biosynthesis by *Crepis rubra*, 307
- biosynthesis and precursors, 307
 - position in triglyceride molecules of seed oil, 307
 - time course of production in seeds, 307
- 7-Octynoic acid, intermediate in unsaturated fatty acid synthesis, 14
- 2-Octynylmalonate, diethyl ester, synthesis, properties, 14
- Oil red O, components staining in serum, 420
- improved lipoprotein staining method, 420
- Oils, marine, cis-9-hexadecenoic and cis-11-octadecenoic acid content of, 313
- Oils, seed, cis-9-hexadecenoic and cis-11-octadecenoic acid content of, 313
- Oils, sunflower, oxygenated acids in, 489
- Oilseeds, *Arctium minus*, trans-3-enoic acids, fatty acid profile, 91
- *Aster alpinus*, trans-2-enoic acids, fatty acid profile, 91
 - *Caltha palustris* L., cis-5-unsaturated fatty acids, 37
 - *Cardamine impatiens* seed oil, triglycerides, 215
 - *Coriaria myrtifolia*, 425
 - *Coriaria nepalensis*, 425
 - *Crepis rubra*, crepenynic acid in triglycerides, 307
 - *Ephedra nevadensis*, fatty acid profile, 170
 - *Ephedra nevadensis*, triglyceride profile, 170
 - *Gossypium hirsutum* L. (cotton plant), carotenoids in plant tissues, 495
 - *Gossypium* sp., lipids of bud, 373
 - *Helianthus annuus* (sunflower), oxygenated acids in, 489
 - *Vernonia anthelmintica*, lipase of, 498

- Olefin**, in crickets, 250
Oleic acid, in bat brown adipose tissue, 340
 — in crickets, 247
 — desaturation to crepenynic acid, 307
 — incorporation into fatty acids of *Crepis rubra*, 307
 — methyl ester, OsO_4 oxidized, mass spectra, 379
 — methyl ester, rate of periodic-permanganate oxidation, 96
 — rabbit liver microsomal phospholipids, variation with age, 361
 — in triglyceride, hydrolysis by *Geotrichum candidum*, 143
 — 12-trimethylsilyloxy, mass spectrum, 431
 — in wheat lipids, maturation changes, 557
Oleic acid (18:1), methyl ester, NMR spectrum, 193
Onosma auriculatum Auch., fatty acid profile, properties, 43
Onosma cinerea Schreb., fatty acid profile, properties, 43
Onosmodium occidentale Mackenz., fatty acid profile, properties, 43
Ophiophagus hannah, phospholipase A, 107
Oxidation, periodic-permanganate, dtmn double bond position, 96
 α -Oxidation, of fatty acids, cofactors required, 275
Oxidation products, of cholesterol, 551
Oxidative cleavage, of unsaturated fatty acids, 37
Oxidation, auto-, of palmitic acid, methyl ester, 59
7-Oxocholesterol trimethylsilyl ethers, mass spectrum, 391
24-Oxocholesterol trimethylsilyl ethers, mass spectrum, 391
Oxygenated acids, HBR-reactive, 459
Ozonolysis, double bond dtmn, 403
Ozonolysis, micro, and GLC of hydroxydienoid fatty esters, 425

P

- Palmitic acid**, in bat brown adipose tissue, 340
 — in crickets, 247
 — ethyl ester, trimethylsilyloxy, mass spectrum, 431
 — metabolism in rat adipose tissue, PGE, effect, 383
 — methyl ester, autooxidation to the monohydroperoxide, 59
 — methyl ester, autooxidation procedure, 65
 — methyl ester, highly purified by preparative GLC, 59
 — methyl ester, hydroperoxide, dtmn of position, 65
 — methyl ester, α -nonadeuterio-trimethylsilyloxy, mass spectrum, 431
 — methyl ester, polarogram of autooxidized product, 59
 — methyl ester, trimethylsilyloxy, mass spectrum, 431
 — methyl ester, 9-trimethylsilyloxy, mass spectrum, 431
 — rabbit liver microsomal phospholipids, variation with age, 361
 — solution with sodium taurocholate, 51
 — trideuteromethyl ester, trimethylsilyloxy, mass spectrum, 431
 — in wheat lipids, maturation changes, 557
Palmitoleic acid, in crickets, 247
 — decrease in phospholipids during pregnancy, 349
 — rabbit liver microsomal phospholipids, variation with age, 361
Pancreas, human, fatty acid methyl esters, 321
 — human, lipids comp, 321
Pancreatic lipase, on dimethyl phosphatides, 191
 — See lipase
Pancreatic lipase EC-3.1.1.3, hydrolysis of triglycerides containing dihydroxy acyl groups, 215
Papain EC-3.4.4.10, effect of lipids, 221
Partition coefficients, carotenes and xanthophylls, 5
Partition number, *Ephedra nevadensis* lipids, 170
n-Partition-2-enal, product of autooxidation, 88
Pepsin EC-3.4.4.1, effect of lipids, 221
Pericardial adipose tissue, dog, triglyceride profiles, 539
Peridinin, of *Gyrodinium resplendens*, 5
 — spectra, ultra-violet, visible; column chromatography, 5
Peridinin (= sulcatoxanthin), spectra, ultra-violet, visible, 5
Peridinol-furanoid-oxide, spectra, ultra-violet, visible; column chromatography, 5
Periplaneta americana. See insects
Perirenal adipose tissue, dog, triglyceride profiles, 539
Perirenal, rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
Periwinkle, triglycerides, positional distribution of fatty acids, 24
Peroxide detection, 59
Peroxides, autooxidation, 331
PGE. See prostaglandin
Phosphatase acid EC 3.1.3.2, effect of lipids, 221
Phosphate, incorporation into sheep erythrocytes *in vitro*, 267
Phosphatidic acid, human brain, equations for computation, 284
 — oxidation during prep, 456
 — phosphate incorporation in sheep erythrocytes, 267
 — TLC, 187
Phosphatidyl acid-like lipid, enzyme inhibition by, 466
Phosphatidyl choline, adsorption by brain particles, 439
 — biosynthesis in liver, variation with age, rabbit, 361
 — in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
 — in *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
 — cotton buds, fatty acid profile and positional distribution, 373
 — fatty acid profile, camel, ass and pig, 107
 — fatty acid profile, goldfish muscle, 121
 — fatty acid profile, mouse brain, 79
 — fatty acid profile, of endometrium, 349
 — from fish, polyunsaturated, ten to twelve double bonds, 191
 — in *Heteropneustes fossilis* (fish) plasma erythrocytes, brain, liver, heart, kidney, 21
 — human brain, equations for computation, 284
 — IR identification, 384
 — IR spectrum, 104
 — in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
 — in mammalian milk, 101
 — molecular species, 107
 — from potato tubers, 477
 — positional distribution of fatty acids, similarity to triglyceride isomers, 30
 — rat brain myelin development, 157
 — rat liver mitochondria, effect of diet on fatty acids, 449
 — TLC, 187
Phosphatidyl choline-cholesterol acyltransferase, serum, 381
Phosphatidyl ethanolamine, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
 — in *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
 — cotton buds, fatty acid profile, 373
 — effect on blood coagulation, 234
 — effect on prothrombin conversion, 234
 — fatty acid profiles, camel, ass and pig, 107
 — fatty acid profile, of endometrium, 349
 — fatty acid profile, mouse brain, 79
 — in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
 — human brain, equations for computation, 284
 — identification, fatty acid profile of *Pseudomonas aeruginosa* grown in hexadecane, 460
 — IR identification, 384
 — IR spectrum, 104
 — in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
 — in liver, variation of fatty acid profile with age, rabbit, 361
 — in mammalian milk, 101
 — metabolism in rat brain, 317
 — molecular species, 107
 — a phosphonic acid analogue of, 234
 — from potato tubers, 477
 — rat brain myelin development, 157
 — rat liver mitochondria, effect of diet on fatty acids, 449
 — TLC, 187
rac-Phosphatidyl (dilinoleyl) ethanolamine, synthesis and properties, 234
Phosphatidyl glycerol, fatty acid profile, in *Spirulina platensis* (algae), 46
 — metabolism in green and blue-green algae, 354
Phosphatidyl hydroxy-L-proline, effect on blood clotting, 228
 — total synthesis of the dioleoyl compound, properties, 228
Phosphatidyl (dioleoyl) hydroxy-L-proline, synthesis and properties, 228
Phosphatidyl inositol, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
 — *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
 — cotton buds, fatty acid profile, 373
 — fatty acid profiles, camel, ass and pig, 107
 — in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
 — human brain, equations for computation, 284
 — *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
 — in mammalian milk, 101
 — from potato tubers, 477
Phosphatidyl methyl transferase, in liver, variation with age, rabbit, 361

- Phosphatidyl phenols**, synthesis from diglycerides, TLC, 519
- Phosphatidyl serine**, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
- *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
- fatty acid profiles, camel, pig and ass, 107
- fatty acid profile, mouse brain, 79
- in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
- human brain, equations for computation, 284
- in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
- in mammalian milk 101
- rat brain myelin development, 157
- TLC, 187
- Phosphoglycerides**, fungus, acetate and glucose incorporation, 211
- Phospholipase A**, *Crotalus atrox* venom, stereospecific anal, 519
- inhibition of alkenyl hydrolase, microsomes, 111
- Phospholipase A₂**, activity in rat liver mitochondria, dependence on diet, 449
- Phospholipase-A EC 3.1.1.4** (Russel viper), on phospholipids, 234
- on polyunsaturated phosphatidyl choline, 191
- stereospecific triglyceride anal, 24
- Phospholipase C**, inhibition of alkenyl hydrolase, microsomes, 111
- Phospholipase C-EC 3.1.4.3**, on polyunsaturated phosphatidyl choline, 191
- Phospholipase D-EC 3.1.4.4**, on polyunsaturated phosphatidyl choline, 191
- Phospholipids**, and accumulation of chemicals, 459
- cholesterol complex, 157
- comp of human liver, 287
- comp of sheep erythrocytes, 267
- distribution in submammalian species, 21
- enzyme activation, 111
- ether, biosynthesis from glyceryl ethers, 51
- fatty acids of, in human erythrocytes, 387
- fatty acid profile, of camel, ass and pig milk, 107
- fatty acid profile, goldfish muscle, 121
- fish, effect of temp, 121
- fungal spores, relation to accumulation of chemicals, 459
- in human pancreas, 321
- incorporation of phosphate into sheep erythrocytes *in vitro*, 267
- influence of sex hormones on EFA deficiency in rats, 199
- mouse brain, fatty acid profile, 79
- mouse brain, fatty aldehyde profile, 79
- in potato tubers, 477
- rat brain myelin development, 157
- rat, tissues, 129
- TLC, 187
- Phospholipids profile**, of mammalian milks, 101
- Photographic reproductions**, of serum lipoproteins, 420
- Photosynthesis**, fatty acid metabolism during, 354
- lipid metabolism during, 354
- Phthalate ester artifacts**, from Negal plastic bottles, 239
- 2-Phthalimidoethyl phosphonic acid**, dicyclohexylammonium salt, synthesis and properties, 234
- synthesis and properties, 234
- Phycomyces**. See microorganisms
- Phytanic acid**, in glycerides and cholesteryl esters, 389
- prep from phytol, TLC, GLC, 389
- 3,7,11,15-tetramethylhexadecanoic acid, 389
- Phytoene**, from cotton plants, properties, 495
- Phytofluene**, from cotton plants, properties, 495
- Pigeon**, *Columba livia intermedia*, phospholipids of tissues, 21
- Pig**, isomer distribution in monoenoic esters, 313
- Pigments**, from potato tubers, 477
- Pig milk**, phospholipids profile, 101
- phospholipids, fatty acid profiles, 107
- Pilchard (South Africa)**, isomer distribution in monoenoic esters, 313
- Plankton**. See microorganisms
- Plant pigments**, fatty acid profile of Xanthophyll esters, 183
- Plants**, *Gossypium hirsutum* L. (cotton plant). See oilseeds
- *Lotus* (clover), lipids attractive to honey bees, 530
- *Melilotus* (clover), lipids attractive to honey bees, 530
- *Solanum tuberosum* (potato). See tubers
- *Tagetes erecta* petals, fatty acid profile of Xanthophyll esters, 183
- *Trifolium* (clover), lipids attractive to honey bees, 530
- Plasma**, dog, 4-¹⁴C-cholesterol turnover, 324
- rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
- Plasma lipids**, rat, influence of sex hormones on EFA deficiency in rats, 199
- Refsum's syndrome, diagnosis by TLC, 389
- Plasmalogen**, fatty acid profiles, camel, ass and pig, 107
- human brain, equations for computation, 284
- microsomal alkenyl hydrolase, 111
- rat brain myelin development, 157
- Plasmalogen**, choline, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
- fatty aldehyde profile, mouse brain, 79
- in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
- Plasmalogen**, ethanolamine, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Columba livia* (pigeon) plasma erythrocytes, brain, liver, heart, kidney, 21
- fatty aldehyde profile, mouse brain, 79
- in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
- Plasmalogen**, serine, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Columba livia* (pigeon), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Heteropneustes fossilis* (fish), plasma erythrocytes, brain, liver, heart, kidney, 21
- in *Kachuga Smithi* (turtle), plasma erythrocytes, brain, liver, heart, kidney, 21
- Polar bear**, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
- triglycerides, positional distribution of docosapentaenoic acid (22:6), 417
- Pollen**, clover (*Melilotus*, *Trifolium*, *Lotus*), lipids attractive to honey bees, 530
- Polyethylene column**, for chromatography, 105
- Polypropylene**, contamination by lipids, 192
- Polystyrene gel**, column chromatography, 385
- Polyunsaturated fatty acids**. See fatty acids, unsaturated, poly-
- Positional distribution**, of fatty acids in triglycerides, 24
- Potato tubers** (*Solanum tuberosum*), lipid components, pigments, 477
- Pregnancy**, effect on fatty acid profiles of phospholipids of endometrium, 349
- Pregn-5-en-3 β -ol**, GLC retention times for trimethylallyl ethers, principal ions in mass spectra, 391
- Pre- and postnatal development**, of rabbit liver lipids, 361
- Pristanic acid**, presence in phytanic acid, 389
- 2,6,10,14-tetramethylpentadecanoic acid, 389
- Pronase**, effect of lipids, 221
- Propanal**, product of autooxidation, 88
- 1,2-Propanediol 3-hexadecyl**. See α -chimy alcohol
- 1,2-Propanediol 3-octadecyl**. See α -batyl alcohol
- 1,3-Propanediol 2-hexadecyl**. See β -chimy alcohol
- 1,3-Propanediol 2-octadecyl**. See β -batyl alcohol
- Prostaglandin**, effect on lipolysis and resynthesis in rat tissues, 383
- effect *in vitro*, on rat adipose tissue, 383
- Protein**, myelin development, 157
- Protein-retinol interaction**, model, 221
- Proteolipids**, elution from dextran gel columns, 453
- Prothrombin conversion**, effect of dioleoylglyceryl(2-aminoethyl)phosphonate, 234
- effect of phosphatidyl ethanolamine (as dioleoyl and dilinoleoyl compounds), 234
- Pseudococcus comstocki** Kuwana, content of dibasic acid, 186
- Pseudomonas aeruginosa**. See microorganisms
- Puccinia graminis (Pers.) f. sp. Tritici**, fatty acid comp, synthesis, 163
- PUFA**. See fatty acids, unsaturated, poly-
- Pyrrhoxanthin**, of *Gyrodinium aureolum*, 5
- spectra, ultra-violet; visible, column chromatography, 5
- Pyrrhoxanthin-furanoid-oxide**, spectra, ultra-violet, visible; column chromatography, 5
- Pyruvic acid kinase EC 2.7.1.40**, effect of lipids, 221

R

- Rabbit**, effect of age and development on liver lipids, 361
- Rabbit endometrium**, phospholipid comp, acetic acid incorporation, 349
- Rabbit serum**, phosphatidyl choline-cholesterol acyltransferase, 381
- Radio isotope**, measurement in tissue of ³H and ¹⁴C, 225
- Rana pipiens**. See frog

- Rat**, adipose tissue, PGE, on glycerol release, 383
 — adipose tissue, PGE, on metabolism, 383
 — adipose tissue, PGE, on palmitic acid metabolism, 383
 — dietary fat effect on fatty acid profile of serum total lipids, 545
 — dietary fat effect on metabolism of acetoacetic acid, 545
 — essential fatty acid deficiency, 199
 — fat deficient, supplemented by experimental lipid comps or linoleic acid, liver lipids, 14
 — fatty acids of skin surface lipid, 403
 — fed corn oil or lard, effect on acetate use, 346
 — Holtzman, hypolipemia, effect of dietary calcium, 1
 — hypolipemia, effect of dietary calcium, 1
 — intestinal contents, metabolism of alkyl glyceryl ethers, 51
 — intestine, washed, metabolism of alkyl glyceryl ethers, 51
 — isomer distribution in monoenoic esters, 313
 — metabolism of alkyl glyceryl ethers, 51
 — Wistar, hypolipemia, effect of dietary calcium, 1
Rat brain, lipid adsorption by particle, 439
 — metabolism, chimyl alcohol, 317
 — metabolism, phosphatidyl ethanolamine, 317
 — microsomal α -oxidation of fatty acids, 275
 — myelin development, 157
Rat intestinal slices, metabolism, effect of diet, 346
Rat kidney, lysosomal cathepsin-D, EC 3.4.4.23, 221
Rat liver, lipid content, effect of strain variety, 1
 — lipids, experimental diets, 14
 — lysosomal cathepsin-D, EC 3.4.4.23, 221
 — metabolism of alkyl glyceryl ethers, 51
 — synthesis, species of diglycerides and triglycerides, 411
 — triglycerides, positional distribution of fatty acids, 30
 — triglycerides, structure, 30
Rat serum, phosphatidyl choline-cholesterol acyltransferase, 381
Rat tissue, lipids, 129
Refsuma's syndrome, diagnosis, chromatographic separation of plasma lipids, 359
Renal. See kidney
Reptiles. See specific name
Retinol, binding to proteins, 221
 — inhibition of enzymes, 221
Retinol-protein interaction, model, 221
Rhizopus nigricans, fungal spore phospholipids, fatty acid profile, 459
Ricinoleic acid, in *Claviceps speciosa*, 260
Ricinoleic acid. See 12-hydroxy-*cis*-9-octadecenoic acid
Rindera lanata (Lam.) Bunge, fatty acid profile, properties, 43
Rindera umbellata (W.K.) Bunge, fatty acid profile, properties, 43
Rochelia disperma (L.) Wettst., fatty acid profile, properties, 43
Rochelia stylaria Boiss., fatty acid profile, properties, 43
Rohrschneider constants, for GLC, 374
Rohrschneider system, for column classification, 374
Royal jelly, fatty acids content, from honey bees, 535
Ruminates, bile acids, comp, 72
- S**
- Salts**, effect on lipid adsorption, 439
Saponification, fatty acids, losses during the extraction of small samples, 189
Scallop, triglycerides, positional distribution of fatty acids, 24
 — triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
Sclerotia, fungal fatty acid profile of *Claviceps* species, 260
Seal, isomer distribution in monoenoic esters, 313
Seal (Alaskan), isomer distribution in monoenoic esters, 313
Sea water, sterol content, 239
Sedimentation velocity, β -apoprotein from β -lipoprotein, 463
Seed lipids, distribution of crepenynic acid in *Crepis rubra*, 307
Seeds, Douglas fir (*Pseudotsuga menziesii* Franco), 482
Sei whale, triglycerides, positional distribution of fatty acids, 24
 — triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
Selachyl alcohol, glyceryl ethers of, identification and quantitation 511
Sephadex chromatography. See chromatography, Sephadex
Serum, blood, in Triton hyperlipemia, dog, 84
 — lipoproteins, α - and β -free and ester cholesterol concentration of, in the baboon, 136
 — lipoproteins α - and β -free, incorporation of ^3H - and ^{14}C -labeled cholesterol and cholesterol esters in the baboon, 136
 — lipoproteins, electrophoresis in a starch gel, 420
Serum lipids, rat, in rat hypolipemia, effect of strain variety, 1
Serum α -1-lipoproteins, human, electrophoretic types, sex differences, photographic reproductions, 420
Serum phosphatidyl choline-cholesterol acyltransferase, effect of Triton-induced hyperlipemia, 381
 — influence of hypercholesterolemia, 381
 — in rat, chicken, rabbit, 381
Sex hormones, effect on essential fatty acid deficiency, rats, 199
Sheep, bile acids, comp, 72
 — erythrocytes, phospholipid comp, 267
Sheep milk, phospholipids profile, 101
Sheepshead (fresh water drum), triglycerides, positional distribution of fatty acids, 24
Sialic acid. See neuraminic acid
Silicones for GLC, phenyl substituted, 374
Silylation, in GLC of triglyceride hydrolysates, 425
 β -Sitosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 — in marine waters. GLC, TLC, 239
Skate (barn-door), triglycerides, positional distribution of fatty acids, 24
Sludge, activated, treatment of sewage, sterol removal, 301
Sodium decyl sulfate, use in solubilization of succinylated apcprotein from β -lipoprotein, 463
Sodium glycocholate, effect on aortic hydrolysis of cholesterol, 454
Sodium taurocholate, effect on aortic hydrolysis of cholesterol, 454
Solanum tuberosum. See tubers
Solubilization of apoprotein, from β -lipoprotein by succinylation, 463
Speckled trout, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
Spectra, gas-liquid-mass, sterol trimethylsilyl ethers, 391
Spectra, infrared, acetoxyhydroxy acid, methyl ester, 215
 — alk-1-enyl ethers, cyclic acetals, 129
 — cholesterol, 551
 — cholesterol-5,6-oxides, 335
 — 3,10-dihydroxydecanoic acid, 535
 — N,N-dimethyl phosphatidyl ethanolamine, 104
 — glucosyl ceramide and galactosyl ceramide, 262
 — 7- α and 7- β -hydroxycholesterol, 551
 — identification of phospholipids, 384
 — 7-ketcholesterol, 551
 — methyl tetradecan-1,14-dioate, 186
 — nitrogenous phospholipids dtmn 9 and 11 microns, 104
 — phosphatidyl choline, 104
 — phosphatidyl ethanolamine, 104
 — of phosphatidyl ethanolamine and N-methyl substituted phosphatidyl ethanolamine, 384
 — of phospholipids, 384
 — of sterols, 239
Spectra, mass, acetoxyhydroxy acid, methyl ester, 215
 — cholesterol-5,6-oxides, 335
 — of di-TMS derivatives of hydroxylated monounsaturated fatty acids, 379
 — hydroxy fatty acids, 431
 — methyl-14-methyl pentadecanoate, 368
 — methyl-12-methyl tetradecanoate, 368
 — methyl-12-methyl tridecanoate, 368
 — oleic acid, 12-trimethylsilyloxy, 431
 — palmitic acid ethyl ester, trimethylsilyloxy, 431
 — palmitic acid methyl ester, α -nonadeuterio trimethylsilyloxy, 431
 — palmitic acid methyl ester, trimethylsilyloxy, 431
 — palmitic acid methyl ester, 9-trimethylsilyloxy, 431
 — palmitic acid trideuteriomethyl ester, trimethylsilyloxy, 431
 — sterol trimethylsilyl ethers, 391
Spectra, mass-gas-liquid chromatography, of di-TMS derivatives of hydroxylated monounsaturated fatty acids, 379
Spectra, nuclear magnetic resonance, acetoxyhydroxy acid, methyl ester, 215
 — aliphatic systems, double bond protons, 193
 — *cis,cis,cis*, 5,11,14,17-eicosatetraenoic acid methyl ester, 37
 — fatty acids, coupling constants, chemical shifts, 193
 — fatty acid methyl esters, 455
Spectra, optical rotary dispersion, 3,10-dihydroxydecanoic acid, 535
Spectra, ultra-violet, inosine, 184
 — visible, β -carotene, 5
 — visible, chlorophyll a, 5
 — visible, chlorophyll c, 5
 — visible, chloroplast pigment Gy 439, 5
 — visible, chloroplast pigment Gy 442, 5
 — visible, chloroplast pigment Gy 443 (a+b), 5
 — visible, diadinoxanthin, 5
 — visible, diadinoxanthin-furanoid-oxide, 5
 — visible, dinoxanthin, 5

- visible, dioxanthin-furanoid-oxide, 5
 - visible, peridin, 5
 - visible, peridin-furanoid-oxide, 5
 - visible, pyrrhoxanthin, 5
 - visible, pyrrhoxanthin-furanoid-oxide, 5
 - Sphingolipid**, biosynthesis in human leucocytes, 151
 - Sphingomyelin**, in *Bufo malanostictus* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
 - in *Columba livia* (pigeon) plasma erythrocytes, brain, liver, heart, kidney, 21
 - fatty acid profiles, camel, ass and pig, 107
 - fatty acid profile, goldfish muscle, 121
 - fatty acid profile, mouse brain, 79
 - in *Heteropneustes fossilis* (toad), plasma erythrocytes, brain, liver, heart, kidney, 21
 - human brain, equations for computation, 284
 - in *Kachuga Smithi* (turtle) plasma erythrocytes, brain, liver, heart, kidney, 21
 - in mammalian milk, 101
 - rat brain myelin development, 157
 - TLC, 187
 - Sphingomyelinosis**, Niemann-Pick disease, 287
 - Spirulina platensis**, fatty acid profiles, total lipids and neutral lipids, 46
 - Spleen**, rat, glyceryl alk-1-enyl ether and glyceryl alkyl ether, 129
 - Squid**, triglycerides, positional distribution of docosahexaenoic acid (22:6), 417
 - Squid**, (see arrow), triglycerides, positional distribution of fatty acids, 24
 - Starvation**. See also fasted animals
 - Stationary phases** for GLC, characterization, 374
 - Stearic acid**, in crickets, 247
 - rabbit liver microsomal phospholipids, variation with age, 361
 - Stereospecific triglyceride analysis**, aquatic animals, 24
 - Sterol glycoside**, cotton buds, fatty acid profile, 373
 - esterified, in potato tubers, 477
 - Sterols**, autooxidation, 239
 - of domestic sewage, 301
 - of house cricket, 256
 - removal by activated sludge, 301
 - trimethylsilyl ethers, 391
 - A5,7-Stigmastadiene-3 β -ol**, in crickets, 256
 - A5,22-Stigmastadiene-3 β -ol**, in crickets, 256
 - Stigmastane-3 β -ol**, in crickets, 256
 - Stigmastanol**, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - A5-Stigmastene-3 β -ol**, in crickets, 256
 - A7-Stigmastene-3 β -ol**, in crickets, 256
 - Stigmasterol**, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391
 - in marine waters, GLC, TLC, 239
 - Storage**, sunflower seeds, changes in oils, 489
 - Stored sunflower seeds**, fatty acid profile of oil from, 489
 - Subcellular distribution**, lipase of germinating fir seeds, 482
 - Subcellular fractions**, of germinating fir seeds, electron microscopy, 482
 - Submammals**. See specific common name
 - Succinylated human serum β -lipoprotein and β -apoprotein**, prep and properties, 463
 - Sulfatase**, aryl EC 3.1.6.1, effect of lipids, 221
 - Sulfatide (cerebroside)**, human brain, equations for computation, 284
 - Sulfatides**. See also cerebroside sulfate
 - Sulfohydriyl enzyme**, involvement in acetylenic fatty acid synthesis in *Crepis rubra*, 307
 - Sulfolipid**. See sulfoquinovosyl diglyceride
 - Sulfoquinovosyl diglyceride**, fatty acid profile, in *Spirulina platensis* (algae), 46
 - metabolism in green and blue-green algae, 354
 - Sunflower oil**, oxygenated acids in, 489
 - Sunflower seeds**, Russian high oil varieties, 489
- T**
- Tagetes erecta**, fatty acid profile of Xanthophyll esters, 183
 - Taurocholic acid**, sodium salt, effect on aortic hydrolysis of cholesteryl oleate, 464
 - Tay-Sachs disease**. See gangliosidosis
 - Temperature**, effect on fatty acid oxidation *in vitro*, 340
 - effect on fatty acid profile of fish, 121
 - Testosterone**, influence on lipids, influence of sex hormones on EFA deficiency in rats, 199
 - Tetradeca-5,8-dienoic acid (14:2 ω 6)**, methyl ester, synthesis, properties, metabolism in rats, 14
 - Tetradeca-5,8-dynoic acid**, synthesis, properties, 14
 - Tetradecan-1,14-dioic acid**, isolation and identification, 186
 - Tetramethyl ammonium hydroxide**, use in extraction of monoenoic acid oxidation products, 96
 - 3,7,11,15-Tetramethylhexadecanoic acid**, phytanic acid, 389
 - 2,6,10,14-Tetramethylpentadecanoic acid**, pristanic acid, 389
 - Thin-layer chromatography**. See chromatography, thin-layer
 - Thiobarbituric acid test**, oxidation products, 331
 - Thyroxine**, lipid antioxidant property, 182
 - Tissue combustion**, for radio isotope dtmn, 225
 - Tissue lipids**, rat, plasma, perirenal, muscle, marrow, liver, spleen, kidney, heart and brain, 129
 - Tissue solubilization**, by hyamine base or NCS, 225
 - TMS (trimethylsilyl)**. See parent compound
 - Toad**, *Bufo malanostictus*, phospholipids of tissues, 21
 - Tobacco budworm**. See insects
 - Tocol**, *in vivo* antioxidant, 331
 - α -Tocopherol**, lipid antioxidant property, 182
 - See 6-chromanol, 2,5,7,8-tetramethyl-(4,8,12-trimethyl tridecyl)
 - α -Tocopherol deficiency**. See also antioxidant deficiency
 - Total lipids**, in liver, variation of fatty acid profile with age, rabbit, 361
 - Trees**, *Pseudotsuga menziesii* Franco (Douglas fir) seeds, 482
 - See also seeds
 - Trichodesma indicum** R. Br., fatty acid profile, properties, 43
 - Trifolium**. See plants
 - Triglycerides**, acetate and glucose incorporation, 211
 - anal of enzymatic hydrolyses by GLC, 425
 - in aquatic animals, 417
 - *Cardamine impatiens* seed oil, acetylated dihydroxy acyl groups, 215
 - comparison of found to calculated profile, 471
 - distribution of oxygenated acyl groups in, 425
 - effects on fatty acid oxidation *in vitro*, 340
 - fatty acid profile, goldfish muscle, 121
 - fatty acid profile, positional distribution, in dog adipose tissue, 539
 - *Geotrichum candidum* lipase, 143
 - GLC, 170
 - human brain, equations for computation, 284
 - in human pancreas, 321
 - hydrolysis by *Vernonia anthelmintica* lipase, 498
 - incorporation of 14 C-glycerol, 411
 - influence of hex hormones on EFA deficiency in rats, 199
 - intramolecular distribution of crepenynic acid, 307
 - isolation from lipid extracts, 30
 - isomers. See triglycerides, positional distribution of fatty acids
 - liquid-liquid partition (reverse phase) chromatography, 170
 - oleic, linoleic and linolenic acids in β -position, 215
 - positional distribution of docosapentaenoic acid and docosahexaenoic acid, 417
 - positional distribution of fatty acids in aquatic animals, 24
 - positional distribution of fatty acids in rat liver, 30
 - positional distribution of fatty acids, similarity to phosphatidyl choline isomers, 30
 - profile, of bovine milk fat, 291
 - profile in human colostrum and milk, 471
 - from potato tubers, 477
 - random utilization of diglycerides in synthesis, 411
 - separation of positional isomers, 30
 - stereospecific anal with *Geotrichum candidum* lipase, 519
 - structure dtmn, 291
 - TLC, GLC, 291
 - Trillineol**, in triglyceride, hydrolysis by *Geotrichum candidum*, 143
 - Trimethylsilyl ethers**, hydroxy fatty acids, GLC, 431
 - sterols, GLC, mass, GLC-mass spectrometry, 391
 - 5,7,8-Trimethyl tocol (α -tocol)**, *in vivo* antioxidant, 331
 - Triolein**, in triglyceride, hydrolysis by *Geotrichum candidum*, 143
 - Triple bond**, introduction into long-chain fatty acid, 307
 - Tritium**, measurement in tissues, 225
 - Triton-induced hyperlipemia**, effect on serum phosphatidyl choline-cholesterol acyltransferase, 381
 - Triton WR-1339**, induction of hyperlipemia, 381
 - Trout (speckled)**, triglycerides, positional distribution of fatty acids, 24
 - Trypsin EC-3.4.4.4**, effect of lipids, 221
 - Tubers**, potato (*Solanum tuberosum*), lipid content, 477
 - *Solanum tuberosum* (white potatoes), lipid content, linoleic acid, linolenic acid, 477
 - Turtle**, *Kachuga Smithi*, phospholipids of tissues, 21
 - Turtle (leatherback)**, triglycerides, positional distribution of fatty acids, 24

U

- UDP-Galactose: N-acetyl galactosaminyl-(N-acetyl neuraminyl) galactosyl-glucosyl-ceramide transferase, in frog brain, 378
 Ultracentrifugally separated lipoproteins, appearance on migration in a starch gel, 420
 Urea, inhibition of alkenyl hydrolase, microsomes, 111
 Uridine diphospho-. See UDP.

V

- Vacuum pump oil, lipid contaminant, control of, 192
 Ventricle, heart. See heart
 Vernonia anthelmintica. See oilseeds
 Very low density lipoproteins, human, migration during starch gel electrophoresis, 420
 Vinyl ethers, TLC, 129
 Violaxanthin, from cotton plants, properties, 495
 Vitamin A, from cotton plants, properties, 495
 Vitamin-E activity, of tocals, 331
 VLDL. See very low density lipoproteins
 — See lipoproteins, very low density

W

- Water, human brain, equations for computation, 284
 Water contamination, by feces, 301
 Wax, scale insect, isolation of a dibasic acid, 186
 Whale, triglycerides, positional distribution of docosapentaenoic acid (22:5), 417
 Whale (Antarctic), isomer distribution in monoenolic esters, 313
 Whale (Icelandic), isomer distribution in monoenolic esters, 313
 Wheat, Kaw, lipid comp, 557
 — maturing, fatty acid profiles of lipids, 557
 — Pawnee, lipid comp, 557
 Wheat plants, rust-infected, fatty acid comp, synthesis, 163

X-Y-Z

- Xanthophylls, fatty acid ester of *Tagetes, erecta*, 183
 — of *Gyrodinium resplendens*, 5
 — partition coefficients, 5
 X-ray diffraction. See also spectra, x-ray
 Zymosterol, GLC retention times for trimethylsilyl ethers, principal ions in mass spectra, 391

